

QUOTIENT SPACES MODULO ALGEBRAIC GROUPS

JÁNOS KOLLÁR

1. INTRODUCTION

In algebraic geometry one often encounters the problem of taking the quotient of a scheme by a group. Despite the frequent occurrence of such problems, there are few general results about the existence of such quotients. These questions come up again and again in the theory of moduli spaces. When we want to classify some type of algebraic objects, say varieties or vector bundles, the classification usually proceeds in two steps.

First we classify the objects with some extra structure added. In classifying varieties we first parametrize embedded varieties in a fixed projective space $\mathbb{P}^N$. In studying vector bundles we first describe them using a basis in the vector space V of global sections.

Second, we have to get rid of the effect of the extra structure. In the case of varieties this usually means dividing out by the automorphism group of $\mathbb{P}^N$. In the case of vector bundles we take the quotient by the automorphism group of the space of sections V .

There have been several attempts to give a general theory of quotients in algebraic geometry.

Mumford's geometric invariant theory is designed to construct quotients by group actions. This approach works very successfully for the quotient problems arising in the theory of vector bundles. Actually, in this area geometric invariant theory solves two problems at once. It predicts the best equivalence relation by which a quotient might exist (a problem with semistable objects), and then it proceeds to construct the quotient.

In the theory of the moduli of varieties, geometric invariant theory has been less successful. It does not seem to predict the correct class of varieties where a good quotient should exist. The existence of the quotient is rather difficult to establish, and it has been done only over $\mathbb{C}$ [Viehweg95].

Artin developed the theory of algebraic spaces in order to tackle more general quotient problems. It was noticed early on that the quotient of a scheme by a finite group is, in general, not a scheme. (The simplest example is smooth of dimension 3). Naively one can think of algebraic spaces as quotients of schemes by finite groups (this is completely correct for normal algebraic spaces). This approach is inconvenient in practice, and the working definition is different [Knutson71].

The theory of algebraic spaces is rather successful in handling general quotient problems. For instance, the quotient of an algebraic space by a flat equivalence relation is again an algebraic space [Artin74]. This implies that if $g : S' \rightarrow S$ is faithfully flat then descent data for algebraic spaces over S' are always effective.

Actions of groups do not produce flat equivalence relations in general, but they are not far from it. Despite this close relationship, it has been rather difficult to prove quotient theorems modulo group actions.

One can not expect the existence of quotients for arbitrary actions. If a group acts on a topological space, a quotient always exists, but it may not be separated. The notion of a group acting “properly” was developed to ensure the separatedness of a quotient. The concept of a proper action is rather easy to translate to algebraic geometry (2.3). Unfortunately, in concrete examples it is frequently hard to check if the group action is proper or not. One difficulty is that being proper is not a local condition in neighborhoods of orbits. Moreover, in considering moduli of vector bundles, the occurring group actions are oftentimes not proper.

The situation is completely different in the theory of polarized varieties, and especially for canonically polarized varieties. The group actions are almost always proper and checking properness is usually the easiest task. Therefore it is reasonable to pay special attention to proper group actions.

The following emerged as a natural problem in this area:

1.1 Conjecture. *Let X be a scheme on which an algebraic group scheme G acts properly. Then the geometric quotient X/G exists (2.7.3).*

1.2 Clarification. In this generality the conjecture definitely fails. G and X must satisfy some technical assumptions. If G/S has varying fiber dimension, then the upper semicontinuity of dimensions shows that a quotient cannot exist in most cases. Therefore, we have to assume that $G \rightarrow S$ is flat or at least universally open.

[Nagata69] constructed examples of Artin rings in characteristic p with a $\mathbb{Z}_p$ -action whose invariant ring is not Noetherian (6.5.1). (The better known constructions of [Nagata59] are not relevant here. In those examples the quotient is an algebraic space of finite type.) To exclude similar cases, we assume that G and X are of finite type over a base scheme.

The conjecture may hold with these assumptions.

1.3 Main Cases. In most of the applications that have been developed so far, one needs this conjecture only in case $G = PGL(n, K)$ for a field K (for moduli spaces over a field) and for $G = PGL(n, \mathbb{Z})$ (for arithmetic moduli questions).

Several moduli problems have been considered where the group in question is not reductive. For instance, the moduli of parabolic bundles appears naturally as a quotient by a parabolic subgroup of GL [Seshadri82]. In many cases the most economical embedding of a variety is to a weighted projective space. Then we need to take the quotient by the automorphisms of a weighted projective space, which is not reductive in general. In the theory of singularities, many of the naturally occurring groups are unipotent [Greuel-Pfister93].

1.4 Known cases. The results of [Artin74] imply (1.1) if the group action is free. For varieties over $\mathbb{C}$ the conjecture was proved by [Popp73], but his proof sheds no light on the algebraic nature of the problem. He used earlier results of [Holman63; Kaup68] asserting that a quotient exists as an analytic space, and then he put an algebraic space structure on it.

For seminormal complex analytic spaces Grauert developed a theory of quotients by rather general equivalence relations; see [Dethloff-Grauert94] for a survey. It would be interesting to work out the analogous theory for algebraic spaces.

A simpler approach to the quotient problem over $\mathbb{C}$ was outlined in [Mumford-Fogarty82, p.172]. In its original form it still used complex analytic quotients, but recently it has been reformulated in terms of algebraic geometry [Keel94; Viehweg95, 9.16]. This method works when the stabilizers of points are reduced. The latter always holds over $\mathbb{C}$, but gives a rather unnatural restriction in general.

The aim of this article is to prove the existence of quotients in two cases.

1.5 Theorem. *Fix an excellent base scheme S . Let G be an affine algebraic group scheme of finite type over S and X a separated algebraic space of finite type over S . Let $m : G \times X \rightarrow X$ be a proper G -action on X . Assume that one of the following conditions is satisfied:*

(1.5.1) *G is a reductive group scheme over S (7.1).*

(1.5.2) *S is the spectrum of a field of positive characteristic.*

Then a geometric quotient $f : X \rightarrow X/G$ exists, and X/G is a separated algebraic space of finite type over S .

1.6 Comments. (1.6.1) There are many examples of moduli problems where the quotient of a scheme by a group is not a scheme. Thus it is natural to consider the quotient problem for algebraic spaces.

(1.6.2) In most of the applications concerning moduli problems, one has to take quotients by $PGL(n, S)$. For these problems, (1.5) provides a satisfactory answer.

(1.6.3) It is somewhat surprising, but I do not see how to use my methods when S is the spectrum of a field of characteristic zero and the group is not reductive. Fortunately, this case has been settled earlier.

The general methods for constructing moduli spaces as geometric quotients imply that (1.5) immediately yields the existence of a slew of moduli spaces over $\text{Spec } \mathbb{Z}$ (cf. [Mumford-Fogarty82; Kollár85]). These results were previously known only in characteristic zero.

The results are most complete for the arithmetic moduli of surfaces of general type. Let $\mathcal{M}_{K^2, \chi}$ denote the moduli functor of canonical models of surfaces of general type with fixed K^2 and $\chi(\mathcal{O})$.

1.7 Theorem. *There is an algebraic space $M_{K^2, \chi}$ over $\text{Spec } \mathbb{Z}$ which coarsely represents the moduli functor $\mathcal{M}_{K^2, \chi}$. $M_{K^2, \chi}$ is separated and of finite type over $\text{Spec } \mathbb{Z}$. $\square$*

For higher dimensional polarized varieties the situation is not so fully understood. Matsusaka's big theorem [Matsusaka72] is still not known in positive characteristic. There are further problems with the locus of ruled varieties (cf. [Kollár85, p.392]).

One case where the latter problem does not occur is the following. Let $\mathcal{M}_P$ denote the moduli functor of smooth, canonically polarized varieties with Hilbert polynomial P .

1.8 Theorem. *There is an algebraic space M_P over $\text{Spec } \mathbb{Z}$ which coarsely represents the moduli functor $\mathcal{M}_P$. M_P is separated and locally of finite type over $\text{Spec } \mathbb{Z}$. $\square$*

1.9 Contents of the Sections. Section 2 reworks the basic foundational properties of geometric quotients in the category of algebraic spaces. The main reason necessitating this is that among algebraic spaces a geometric quotient need not be unique (2.18). Uniqueness holds for proper actions, but it needs a careful proof.

In section 3 I study general properties of quotient morphisms. The main result (3.12) asserts that if a morphism looks like a quotient topologically, then a geometric quotient exists with the same underlying topological space. These results are related to [Fogarty83].

Next I study the quotient problem for successively larger classes of algebraic spaces.

For normal spaces the quotient problem was considered in [Seshadri63,71]. He did not formulate the results in terms of algebraic spaces, but in essence he proved (1.1) for X normal. These theorems are recalled in section 4.

If X is reduced, we would like to construct the quotient using the normalization $\bar{X} \rightarrow X$. We already know that $\bar{X}/G$ exists, and we just have to undo the effects of normalization. These types of problems have been considered by [Artin70,6.3]. The problem turns out to be rather delicate. For technical reasons we assume that X is weakly normal (5.4). The group theoretic aspects are discussed in section 5; most of the technical problems are left to section 8.

For an arbitrary algebraic space the quotient is built using the already constructed quotient of the weak normalization of X . In positive characteristic the Frobenius morphism can be exploited to give the general result. This is discussed in section 6.

Reductive group schemes are considered in section 7. I have to use the theorems of [Seshadri77] on reductive group schemes acting on affine schemes, though this goes against the general philosophy of my approach.

1.10 Technical Assumptions. Throughout the paper, the base scheme is denoted by S . We always assume that S is locally Noetherian. Starting with section 4, we also assume that it is excellent. This assumption could be weakened a little, but there are serious difficulties arising for general Noetherian base.

Starting with section 3, all algebraic spaces are assumed separated and locally of finite type over S . Lack of separation causes serious problems, but they probably can be overcome through careful examination of the proofs and definitions. In considering group actions, the interesting part always seems to happen in a finite type extension, though I do not know any general result to this effect. The assumption is also important technically.

All group schemes are assumed to be separated and of finite type over S .

Acknowledgements. Partial financial support was provided by the NSF under grant numbers DMS-8707320 and DMS-9102866.

These notes were typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\TeX}$, the $\text{\TeX}$ macro system of the American Mathematical Society.

2. GEOMETRIC QUOTIENTS OF ALGEBRAIC SPACES

The aim of this section is to review the basic definitions and theorems about group actions and various quotients. Most of the material follows [Mumford-Fogarty82, chapter 0] rather closely, reinterpreted in the context of algebraic spaces. Most definitions and results go through without any change. The main reason of repeating the material for algebraic spaces is that there is one subtle and surprising difference:

Geometric quotients are not necessarily categorical quotients any more, and there may exist more than one geometric quotient (2.18).

For proper group actions these anomalies do not occur, but this fact requires a careful proof (2.15).

2.1 Definition. (2.1.1) A *group scheme* over S is a scheme G/S together with morphisms

multiplication: $\mu : G \times_S G \rightarrow G$, inverse: $b : G \rightarrow G$, and unit: $e : S \rightarrow G$, which satisfy the obvious group laws in the scheme theoretic setting (see [Mumford-Fogarty82, 0.1]).

(2.1.2) Let G/S be a group scheme and X/S an algebraic space. An *action of G on X* is a morphism $m_X : G \times_S X \rightarrow X$ such that $m(\mu(p_1, p_2), p_3) = m_X(p_1, m_X(p_2, p_3))$ where the p_i are the coordinate projections of $G \times_S G \times_S X$. We usually say that X is a *G -space* if the action is understood.

If $\psi : U \rightarrow G$ and $f : U \rightarrow X$ are morphisms then $m_X(\psi, f) : U \rightarrow X$ is usually denoted by $\psi \cdot f$.

The *trivial G action* on X is given by $m_X = p_2$ where $p_2 : G \times_S X \rightarrow X$ is the second projection.

(2.1.3) Let G/S be a group scheme, X/S a G -space and F a sheaf on X . An *action of G on F* is an isomorphism $\sigma : m_X^* F \cong p_2^* F$ which satisfies the scheme theoretic analog of the cocycle condition; see [Mumford-Fogarty82, 1.3] for details.

2.1.4 Remark. One can also consider group objects in the category of algebraic spaces, but these never seem to have acquired a good name. Aside from this terminological problem, we could always use this more general notion.

2.2 Definition. (2.2.1) Let $m_i : G \times X_i \rightarrow X_i$ be G -spaces. A morphism $f : X_1 \rightarrow X_2$ is called a *G -morphism* if the following diagram is commutative:

$$\begin{array}{ccc} G \times X_1 & \xrightarrow{m_1} & X_1 \\ id_G \times f \downarrow & & \downarrow f \\ G \times X_2 & \xrightarrow{m_2} & X_2. \end{array}$$

If F is a G -sheaf on X_1 then the direct image $f_* F$ is G -sheaf on X_2 .

(2.2.2) A *G -map of G -sheaves* is defined in the obvious way. If X is an algebraic space with trivial G -action and F a (quasi coherent) G -sheaf on X then

$$X \supset U \mapsto H^0(U, F|_U)^G$$

is a (quasi coherent) sheaf on X . It is denoted by F^G . (The definition makes sense even if the G -action on X is not trivial, but then F^G is not a sheaf of $\mathcal{O}_X$ -modules.)

2.3 Definition. Let X be a G -space. Let Ψ_X denote the morphism

$$\Psi_X : G \times_S X \xrightarrow{(m_X, p_2)} X \times_S X,$$

where $p_2 : G \times_S X \rightarrow X$ is the second projection.

We say that the action of G on X is *proper* if Ψ_X is proper.

2.3.3 Remark. It is somewhat unfortunate that the name “proper” is used to describe a condition which guarantees the separatedness of the quotient (2.9).

The following is a reformulation of the valuative criterion of properness:

2.4 Lemma. *Let X be a G -space. The G action on X is proper iff the following condition is satisfied:*

Let T be the spectrum of a DVR with generic point t_g and $u_1, u_2 : T \rightarrow X$ two morphisms. Assume that there is a morphism $\psi_g : t_g \rightarrow G$ such that $u_1(t_g) = \psi_g(t_g) \cdot u_2(t_g)$. Then there is unique extension of ψ_g to a morphism $\psi : T \rightarrow G$ such that $u_1 = \psi \cdot u_2$. $\square$

2.5 Proposition. *Let $f : X \rightarrow Y$ be a separated G -morphism of G -spaces. If G acts properly on Y then G acts properly on X .*

Proof. By assumption $\Psi_Y : G \times Y \rightarrow Y \times Y$ is proper, By base change we obtain that $\Psi_Y \times_Y X : G \times X \rightarrow Y \times X$ is also proper. The latter can be factored as

$$\Psi_Y \times_Y X : G \times X \xrightarrow{\Psi_X} X \times X \xrightarrow{(f, id_X)} Y \times X.$$

Thus Ψ_X is proper. $\square$

2.6 Definition. Let $f_i : X_i \rightarrow Y$ be G -morphisms of G -spaces. Let $\Delta : G \rightarrow G \times G$ be the diagonal imbedding. Then the fiber product $X_1 \times_Y X_2$ is a G -space via the morphism

$$m : G \times (X_1 \times X_2) \xrightarrow{(\Delta, id)} G \times G \times (X_1 \times X_2) \cong (G \times X_1) \times (G \times X_2) \xrightarrow{(m_1, m_2)} X_1 \times X_2.$$

The coordinate projections $X_1 \times_Y X_2 \rightarrow X_i$ are G -morphisms.

There are various ways of defining the quotient of an algebraic space by a group. The following are the basic variants. I am mostly interested in proper actions, therefore the concept of “good quotient” is not relevant.

2.7 Definition. Let G/S be a group scheme and X/S a G -space. Let Z be an algebraic space with trivial G -action and $q : X \rightarrow Z$ a G -morphism.

(2.7.1) $q : X \rightarrow Z$ is called a *categorical quotient* of X by G if, for every algebraic space U with trivial G -action and G -morphism $p : X \rightarrow U$, there is a morphism $u : Z \rightarrow U$ such that $p = u \circ q$.

(2.7.2) $q : X \rightarrow Z$ is called a *topological quotient* of X by G if the following conditions are satisfied:

(2.7.2.1) f is a G -morphism.

(2.7.2.2) f is locally of finite type.

(2.7.2.3) If K is an algebraically closed field then $f(K) : X(K)/G(K) \rightarrow Z(K)$ is an isomorphism (of sets).

(2.7.2.4) f is universally submersive.

(2.7.3) $q : X \rightarrow Z$ is called a *geometric quotient* of X by G if it is a topological quotient and in addition

$$(2.7.2.5) \mathcal{O}_Z = (q_* \mathcal{O}_X)^G.$$

2.8 Remarks. (2.8.1) Strictly speaking, the definitions (2.2–3) make sense for algebraic spaces only if we establish their invariance under étale morphisms $Z' \rightarrow Z$. This is done in (2.10).

(2.8.2) By [Mumford-Fogarty82, 0.1], in the category of *schemes* a geometric quotient is also a categorical quotient. In particular, it is unique up to isomorphism. We see in (2.18) that for algebraic spaces both of these can fail if the G -action is not proper.

(2.8.3) This definition differs slightly from the one in [Mumford-Fogarty82, sec. 0.1]. I assume that f is locally of finite type. In keeping with the original definition in [Mumford65, sec. 0.1], I assume that f is universally submersive (and not just submersive). As [Mumford65, sec. 0.1] argues, universally submersive is a more topological notion. If G is open over S then the two notions are equivalent by [Dixmier-Raynaud81, 1.9], and it is convenient to have the stronger variant from the beginning. For topological quotients it is crucial to assume universal submersiveness, see (3.12.4).

(2.8.4) By [Mumford-Fogarty82, p.6], if q is a topological quotient and G/S is universally open then q is also universally open.

2.9 Proposition. *Assume that the G -action on X is proper and a topological quotient $f : X \rightarrow Z$ exists. Then Z is separated.*

Proof. f is universally submersive, thus, if T is the spectrum of a DVR and $u : T \rightarrow Z$ a morphism then there is a dominant morphism $h : T' \rightarrow T$ (where T' is the spectrum of a DVR) such that $u \circ h : T' \rightarrow Z$ can be lifted to a morphism $\bar{u} : T' \rightarrow X$ (cf. (3.7)).

Let $u_1, u_2 : T \rightarrow X/G$ be two morphisms which agree at the generic point. We find $T' \rightarrow T$ such that both of them lift to $\bar{u}_i : T' \rightarrow X$. $\bar{u}_1|_{t'_g}$ and $\bar{u}_2|_{t'_g}$ lift the same morphism $t'_g \rightarrow Z$, thus they differ by translation by G . Since the G action on X is proper, the two morphisms $\bar{u}_i$ differ only by a translation by G . Thus $f \circ \bar{u}_1 = f \circ \bar{u}_2$, hence Z is separated. $\square$

2.10 Proposition. [Mumford-Fogarty82, p.9] *Let X be a G -scheme and $f : X \rightarrow Z$ a G -morphism (with the trivial G -action on Z). Let $h : Z' \rightarrow Z$ be a flat morphism. Set $X' := X \times_Z Z'$ and let $f' : X' \rightarrow Z'$ be the second projection.*

(2.10.1) *If $f : X \rightarrow Z$ is a geometric (resp. topological) quotient of $X \bmod G$, then $f' : X' \rightarrow Z'$ is also a geometric (resp. topological) quotient of $X' \bmod G$.*

(2.10.2) *Assume that h is faithfully flat. Then $f' : X' \rightarrow Z'$ is a geometric (resp. topological) quotient of $X' \bmod G$ iff $f : X \rightarrow Z$ is a geometric (resp. topological) quotient of $X \bmod G$. $\square$*

If $f : X \rightarrow Y$ is an étale G -morphism then the induced morphism between the geometric quotients is usually not étale. The following condition was introduced by Deligne (cf. [Knutson71, p. 183]) to remedy this problem.

2.11 Definition. Let X be a G -space, and Ψ_X as in (2.3). The fiber $\Psi_X^{-1}(x, x) \subset G \times \{x\}$ is a subgroup scheme. It is called the *stabilizer* of x , and is denoted by $\text{Stab}(x)$.

2.12 Definition. Let $f : X \rightarrow Y$ be a G -morphism. We say that f is *fixed-point reflecting* if the following condition is satisfied:

$$\text{Stab}(x) = \text{Stab}(y) \times_y \{x\} \text{ for any } x \in X \text{ and } y = f(x).$$

The simplest examples of fixed-point reflecting morphisms are obtained by base change from trivial G -actions:

2.13 Lemma. *Let $f : X \rightarrow Z$ be a G -morphism with trivial G -action on Z and $h : Z' \rightarrow Z$ any morphism. Then $X \times_Z Z' \rightarrow X$ is fixed-point reflecting. $\square$*

The following result shows the importance of fixed-point reflecting morphisms. It is related to [Luna73, Chap. II]. The theorem is also crucial in establishing basic properties of geometric quotients in the category of algebraic spaces.

2.14 Theorem. *Assume that $G \rightarrow S$ is universally open. Let $f : X \rightarrow Y$ be a separated and fixed-point reflecting G -morphism. Assume that the G -action on Y is proper and that f is étale. Assume furthermore that a topological quotient $p_Y : Y \rightarrow Z(Y)$ exists.*

Then there is a topological quotient $p_X : X \rightarrow Z(X)$ such that the induced morphism $Z(X) \rightarrow Z(Y)$ is étale.

Proof. By (2.8.4) p_Y is universally open and surjective. Thus by [SGA1, IX.4.9] all descent data on separated and étale morphisms to Y are effective. (See [BLR90, chapter 6] for a clear and selfcontained discussion on descent.) Therefore it is sufficient to construct a descent datum for $f : X \rightarrow Y$, that is, an isomorphism

$$\phi : X \times_{Z(Y)} Y \xrightarrow{\cong} Y \times_{Z(Y)} X \quad \text{over} \quad Y \times_{Z(Y)} Y$$

which satisfies the cocycle condition [BLR90, p.133].

The construction of ϕ , and the role of the fixed-point reflecting conditions, is transparent at the set theoretic level.

Let K be an algebraically closed field. Pick $x \in X(K)$ and $y \in Y(K)$ such that $p_Y(f(x)) = p_Y(y)$. By (2.7.2.3) there is a $g \in G(K)$ such that $f(x) = g \cdot y$. Set

$$(2.14.1) \quad \phi(x, y) = (g \cdot y, g^{-1} \cdot x).$$

g is not uniquely defined by the equation $f(x) = g \cdot y$; two choices differ by an element of $\text{Stab}(y)$. Since $\text{Stab}(x) = \text{Stab}(y)$, the product $g^{-1} \cdot x$ is independent of the choice of g . Thus ϕ is well defined.

We construct ϕ by constructing its graph. (2.14.1) shows that, at least set theoretically, the graph of ϕ is the image of the morphism

$$\begin{aligned} h : G \times_{Z(Y)} X &\rightarrow V := X \times_{Z(Y)} Y \times_{Y \times_{Z(Y)} Y} Y \times_{Z(Y)} X, \quad \text{given by} \\ h : (g, x) &\mapsto (g \cdot x, f(x), g \cdot f(x), x). \end{aligned}$$

$G \times_{Z(Y)} Y \rightarrow Y \times_{Z(Y)} Y$ is proper, hence so is $h_2 : G \times_{Z(Y)} X \rightarrow Y \times_{Z(Y)} X$. h is the fiber product of two such morphisms, hence h is also proper. Thus $W := \text{im } h \subset V$ is a closed subscheme.

h_2 can be factored as

$$h_2 : G \times_{Z(Y)} X \rightarrow W \xrightarrow{\pi_2} Y \times_{Z(Y)} X.$$

This shows that π_2 is proper. The set theoretic computation of ϕ shows that π_2 is universally injective and surjective, thus π_2 is a universal homeomorphism.

$X \times_{Z(Y)} Y$ and $Y \times_{Z(Y)} X$ are both étale over $Y \times_{Z(Y)} Y$, thus the projection morphisms

$$\begin{array}{ccc} V & \longrightarrow & Y \times_{Z(Y)} X \\ \downarrow & & \downarrow \text{id}_Y \times f \\ X \times_{Z(Y)} Y & \xrightarrow{f \times \text{id}_Y} & Y \times_{Z(Y)} Y \end{array}$$

are all étale.

Thus $W \subset V$ provides a section of $V \rightarrow Y \times_{Z(Y)} Y$, at least after base change to W . Since $W \rightarrow Y \times_{Z(Y)} Y$ is a universal homeomorphism, (5.2) implies that

$V \rightarrow Y \times_{Z(Y)} Y$ has a unique section $W^0 \subset V$ such that $\text{Supp } W^0 = \text{Supp } W$. W^0 is the graph of ϕ .

By (5.3) it is sufficient to check the cocycle condition set theoretically, where it is trivial. $\square$

2.14.2 Remark. (2.14) implies that a fixed-point reflecting and étale G -morphism is also strongly étale in the sense of [Mumford-Fogarty82, p. 152].

2.15 Corollary. *Let X be an algebraic G -space. Assume that the G -action on X is proper. Let $q : X \rightarrow Z$ be a geometric quotient. Then q is also a categorical quotient. Thus a geometric quotient is unique (up to a canonical isomorphism).*

Proof. [Mumford-Fogarty82, p.4] shows that in the category of *schemes* a geometric quotient is also a categorical quotient. More generally, the proof shows the following:

2.15.1 Claim. *Let X be an algebraic G -space and $q : X \rightarrow Z$ a geometric quotient. Let V be a scheme with trivial G -action and $p : X \rightarrow V$ a G -morphism. Then there is a morphism $v : Z \rightarrow V$ such that $p = v \circ q$. $\square$*

In order to get the general case, let U be an algebraic space with trivial G -action and $p : X \rightarrow U$ a G -morphism. We need to find $u : Z \rightarrow U$. To do this let V be an affine scheme and $j : V \rightarrow U$ an étale morphism. By base change we obtain a fixed-point reflecting étale morphism $p^*j : V \times_U X \rightarrow X$.

Applying (2.14) we obtain that a geometric quotient $p' : V \times_U X \rightarrow Z'$ exists and the induced morphism $Z' \rightarrow Z$ is étale. Since V is a scheme, by (2.15.1) we obtain a morphism $Z' \rightarrow V$. The resulting diagram of étale morphisms

$$Z \leftarrow Z' \rightarrow V \rightarrow U$$

gives the local description of $u : Z \rightarrow U$. $\square$

2.16 Notation. Let X be an algebraic G -space such that the G -action on X is proper. The unique geometric quotient is denoted by $q : X \rightarrow X/G$.

2.17 Corollary. *Let $f : X \rightarrow Y$ be a separated, surjective, étale and fixed-point reflecting G -morphism. Assume that the G -action on Y is proper. Assume furthermore that the geometric quotient $q_X : X \rightarrow X/G$ exists.*

Then the geometric quotient $q_Y : Y \rightarrow Y/G$ exists and the induced morphism $X/G \rightarrow Y/G$ is étale.

Proof. Let $Z = X \times_Y X$. Z can be viewed as an étale equivalence relation on X and the quotient of X by this equivalence relation is Y .

Apply (2.14) to $Z \rightarrow X$. We conclude that Z/G exists and the maps $Z/G \rightarrow X/G$ are étale. Thus Z/G gives an étale equivalence relation on X/G . The quotient is an algebraic space U such that $Y \times_U (X/G) \cong X$. Thus by (2.10.2) U is the geometric quotient of $Y \bmod G$. $\square$

The following example, taken from [Mumford-Fogarty82, p.11] shows that a geometric quotient need not be a categorical quotient in general.

2.18 Example. Let $SL(2, \mathbb{C})$ act on homogeneous polynomials by the rule

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} : x \mapsto ax + cy, \quad y \mapsto bx + dy.$$

Let X be the variety of pairs (L, Q^2) and Y the variety of pairs (L, Q) where L is a nonzero linear form and Q is a quadratic form of discriminant -1. It is easy to see that X and Y are smooth, irreducible and $SL(2, \mathbb{C})$ acts freely on them. The morphism

$$f : Y \rightarrow X \quad \text{given by} \quad (L, Q) \mapsto (L, Q^2)$$

is an étale and fixed-point reflecting $SL(2, \mathbb{C})$ -morphism. f has degree two and the fiber over (L, Q^2) has two points (L, Q) and $(L, -Q)$.

A geometric quotient of X is obtained as follows. Every pair can be brought to the form (x, Q^2) . Then

$$X \rightarrow \mathbb{C} \quad \text{given by} \quad (x, Q^2) \mapsto Q(0, 1)^2$$

is a geometric quotient.

Since $\mathbb{C}$ has no nontrivial étale covers, this already shows that (2.14) can fail if the group action is not proper.

The stabilizer of x is the group of upper triangular matrices

$$\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}.$$

On quadratic forms the action of upper triangular matrices is

$$ux^2 + vxy + wy^2 \mapsto (u + vb + wb^2)x^2 + (v + 2wb)xy + wy^2.$$

This shows that (L, Q) and $(L, -Q)$ are always in different orbits.

A geometric quotient of Y is obtained as follows. Every pair can be brought to the form (x, Q) . Then

$$X \rightarrow Z \cong \mathbb{C} \quad \text{given by} \quad (x, Q) \mapsto Q(0, 1)$$

is a geometric quotient, provided $Q(0, 1) \neq 0$. If $Q(0, 1) = 0$ then we get two different orbits corresponding to the pairs (x, xy) and $(x, -xy)$. The geometric quotient is not separated, as shown by the families of pairs

$$(x, xy + ty^2) \quad \text{and} \quad (x, -xy + ty^2).$$

These lie in the same orbit for $t \neq 0$ but in different orbits if $t = 0$. Thus a geometric quotient of Y by $SL(2, \mathbb{C})$ is the nonseparated line (which is a scheme)

$$U : \mathbb{C} \times \{0, 1\} \quad \text{modulo the equivalence} \quad (t, 0) \sim_1 (t, 1) \quad \text{for } t \neq 0.$$

The étale equivalence relation $(t, 0) \sim_2 (-t, 1)$ on $\mathbb{C} \times \{0, 1\}$ descends to an étale equivalence relation on U . Let V denote the quotient (which is an algebraic space).

The morphism

$$\mathbb{C} \times \{0, 1\} \rightarrow Z \cong \mathbb{C} \quad \text{given by} \quad (t, 0) \mapsto t^2, \quad (t, 1) \mapsto t^2$$

descends to a morphism $p : V \rightarrow Z$. p is an isomorphism on the set of points, but it is not an isomorphism. Indeed, $\partial/\partial t$ gives a nonzero tangent vector of V

at the origin, but it maps to zero on Z . V is the simplest example of a locally nonseparated algebraic space (cf. [Knutson71, p.9]).

Finally, $Y \rightarrow U$ descends to an $SL(2, \mathbb{C})$ -morphism $X \rightarrow V$, but there is no corresponding morphism from Z to V . Thus the geometric quotient is not a categorical quotient in the category of algebraic spaces.

One can see that in this case V is both a geometric and a categorical quotient of X .

2.19 Example. Let $S = \mathbb{C}$ and $G_1 := S \times \{+1, -1\}$ the constant 2-element group scheme over S . Let $X \subset \mathbb{C} \times S$ be the union of the diagonal (x, x) and of the anti diagonal $(-x, x)$. G_1 acts on X by multiplication on the first coordinate.

Let $G \subset G_1$ be the subgroup scheme $S \times \{+1\} \cup (S \setminus \{0\}) \times \{-1\}$. G acts on X , the action is closed but not proper and the geometric quotient is S . $X \rightarrow S$ is affine; this is of interest in connection with (3.12).

The G -action on X does not identify the two directions of the tangent cone of X . One can see that the categorical quotient (in the category of algebraic spaces) is not locally separated above $0 \in S$.

2.20 Remark. It is quite likely that a geometric quotient is also a categorical quotient in the category of locally separated algebraic spaces. If $S = \mathbb{C}$ and G is reductive, this follows from the fact that a geometric quotient is also a categorical quotient in the category of complex analytic spaces [Neeman89, Thm. 10].

3. TOPOLOGICAL QUOTIENTS

Let X be a G -space. In this section we assume that we already have a quotient morphism $f : X \rightarrow Z$, and concentrate on analyzing the relationships among its properties. The main result is (3.12), which asserts that a quotient with good topological properties has very good scheme theoretic properties as well.

3.1 Assumptions. Fix a locally Noetherian base scheme S . For the rest of this section we assume that G is locally of finite type over S and the structure morphism $G \rightarrow S$ is universally open.

All algebraic spaces are assumed to be separated. All group actions are assumed to be proper.

3.2 Definition. Let X be a G -space. A morphism $f : X \rightarrow Z$ is called a *weak topological quotient* if the following conditions are satisfied:

- (3.2.1) f is a G -morphism (with the trivial G -action on Z).
- (3.2.2) f is locally of finite type.
- (3.2.3) If K is an algebraically closed field then $f(K) : X(K)/G(K) \rightarrow Z(K)$ is an isomorphism (of sets).

If Z is normal then these properties are quite strong (3.9). The following example shows what can go wrong in general:

3.3 Examples. Let $S = \text{Spec } \mathbb{C}$ and $G = S$ the trivial group scheme.

(3.3.1) Let $Z = \text{Spec } \mathbb{C}[x]$, $X = \{0\} \cup (Z \setminus \{0\})$ (disjoint union) and $f : X \rightarrow Z$ the natural morphism. The conditions (3.2.1–3) are satisfied. A more sophisticated version of this is example is the following.

(3.3.2) Let Z be an irreducible curve with a single node $z \in Z$ and $f' : X' \rightarrow Z$ the normalization with $\pi, \pi' \in X'$ the preimages of z . Set $Y := X' \setminus \pi$ and

$f : X \rightarrow Z$ the restriction of f' . Then f is a homeomorphism and the conditions (3.2.1–3) are satisfied.

Two bad properties of f come to mind first: f is not proper and not flat. Unfortunately, in most cases, the quotient morphism is neither flat nor proper.

The correct approach seems to find suitable variants of the valuative criterion of properness. The next two definitions give first a weak variant and then a much stronger one. At the end, they turn out to be equivalent (3.7).

3.4 Definition. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces. We say that f satisfies the *weak lifting property for DVR's* if the following holds:

Let T be the spectrum of a DVR and $u : T \rightarrow Y$ a morphism. Then there is a commutative diagram

$$\begin{array}{ccc} T' & \xrightarrow{u'} & X \\ h \downarrow & & \downarrow f \\ T & \xrightarrow{u} & Y, \end{array}$$

where T' is the spectrum of a DVR and $h : T' \rightarrow T$ a surjective morphism.

3.5 Definition. (cf. [Seshadri72, 4.1]) Let $f : X \rightarrow Y$ be a G -morphism of G -spaces such that the G -action on Y is trivial.

We say that f is *complete mod G* if f is separated, of finite type and the following version of the valuative criterion of properness is satisfied:

Let T be the spectrum of a DVR with generic point $j : t_g \rightarrow T$. Assume that we have a commutative diagram

$$(3.5.1) \quad \begin{array}{ccc} t_g & \xrightarrow{u_g} & X \\ j \downarrow & & \downarrow f \\ T & \longrightarrow & Y \end{array}$$

Then there is a commutative diagram

$$(3.5.2) \quad \begin{array}{ccc} T' & \xrightarrow{u'} & X \\ h \downarrow & & \downarrow f \\ T & \longrightarrow & Y \end{array} \quad \text{and} \quad \begin{array}{c} \text{a morphism} \\ \phi'_g : t'_g \rightarrow G, \end{array}$$

where T' is the spectrum of a DVR with generic point t'_g and $h : T' \rightarrow T$ a dominant morphism, such that

$$u_g \circ (h|t'_g) = \phi'_g \cdot (u'|t'_g).$$

Informally speaking, the valuative criterion of properness is satisfied after a base change and a suitable translation by the group.

3.5.3 Remarks. (3.5.3.1) If the G -action on X is trivial then f is proper iff it is complete mod G .

(3.5.3.2) Since f is locally of finite type, we may assume in addition that $h : T' \rightarrow T$ is finite.

(3.5.3.3) It is possible to define the notion of “complete mod G ” for G -morphisms where we do not assume that the G -action on Y is trivial. I do not see any advantages at the moment.

The following proposition lists some easy results about the composition of morphisms which are proper or complete mod G . I list only those variants which are used later. The proofs are straightforward.

3.6 Proposition. *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be separated G -morphisms of G -spaces.*

(3.6.1) *If f is proper and g is complete mod G then $g \circ f$ is complete mod G .*

(3.6.2) *If f is complete mod G and g is proper then $g \circ f$ is complete mod G .*

(3.6.3) *If $g \circ f$ is complete mod G and the G action on Y is trivial then f is complete mod G .*

(3.6.4) *If $g \circ f$ is complete mod G , the G action on Y is trivial and f is surjective then g is proper. $\square$*

The next result shows that several good properties of weak topological quotients are equivalent:

3.7 Proposition. *Let X be a G -space and $f : X \rightarrow Z$ a G -morphism, locally of finite type (with trivial G -action on Z). Then each of the following conditions implies the next:*

(3.7.1) *f is surjective and complete mod G .*

(3.7.2) *f is surjective and universally open.*

(3.7.3) *f is universally submersive.*

(3.7.4) *f satisfies the weak lifting property for DVR's.*

If, in addition, $f : X \rightarrow Z$ is a weak topological quotient, then all four conditions are equivalent.

Proof. Assume (3.7.1). (3.7.1) is stable under any base change, thus it is sufficient to prove that f is open. Let $U \subset X$ be open and $V = f(U) \subset Y$. V is constructible by Chevalley's theorem [EGA71, I.7.1.4], hence V is open iff it is closed under generalization.

Pick $y \in V$ and let y' be a generalization of y . Let $h : T \rightarrow Y$ be a morphism such that y' is the image of the generic point and y the image of the closed point. f is surjective, hence there are $x, x' \in X$ such that $f(x) = y$ and $f(x') = y'$. By (3.5) there is a commutative diagram

$$\begin{array}{ccc} T' & \xrightarrow{u'} & X \\ h \downarrow & & \downarrow f \\ T & \xrightarrow{u} & Y. \end{array}$$

By enlarging T' we may assume that T' has algebraically closed residue field K and that x is a K -valued point of X . Thus there is a $\phi_0 : K \rightarrow G$ such that $x = \phi_0 \cdot (u'|_{t'_0})$. After replacing T' by a ramified extension, we may also assume that ϕ_0 extends to $\phi : T' \rightarrow G$ (here we use that G/S is universally open). Set $u'' := \phi \cdot u'$.

$u'' : T' \rightarrow X$ is a morphism such that $x = u''(t'_0)$ and $x'' := u''(t'_g)$ is a generalization of x . Thus $x'' \in U$ and therefore $u'' = f(x'') \in V$. Thus f is open.

(3.7.2) $\Rightarrow$ (3.7.3) holds for any continuous map of topological spaces.

In order to see (3.7.3) $\Rightarrow$ (3.7.4) let $u : T \rightarrow Z$ be a morphism and $t_0 \in T$ the closed point. By assumption $u^*f : X \times_Z T \rightarrow T$ is submersive, thus $(u^*f)^{-1}(t_0) \subset X \times_Z T$ is not open. Thus there is a point $x \in (u^*f)^{-1}(t_0)$ which is the specialization of a point $x' \notin (u^*f)^{-1}(t_0)$. This gives $v : T' \rightarrow X \times_Z T$ such that x' is the image of the generic point and x the image of the closed point. Thus $h : T' \rightarrow T$ is dominant and $u' : T' \rightarrow X \times_Z T \rightarrow X$ gives a lifting.

Finally, assume (3.7.4) and that f is a weak topological quotient. Consider a commutative diagram

$$\begin{array}{ccc} t_g & \xrightarrow{u_g} & X \\ j \downarrow & & \downarrow f \\ T & \xrightarrow{u} & Y. \end{array}$$

By assumption there is a commutative diagram as in (3.4)

$$(3.7.5) \quad \begin{array}{ccc} T' & \xrightarrow{u'} & X \\ h \downarrow & & \downarrow f \\ T & \xrightarrow{u} & Y. \end{array}$$

By enlarging T' we may assume that the quotient field K of T' is algebraically closed. We show that (3.7.5) is the diagram whose existence is required in (3.5.2).

$u'|t'_g$ and u_g are K -valued points of X and they have the same image in Y . Since f is a weak topological quotient, there is a $\phi'_g : t'_g \rightarrow G$ such that $u_g \circ (h|t'_g) = \phi'_g \cdot (u'|t'_g)$. This shows (3.7.1). $\square$

3.7.6 Remark. The above proof shows that if $f : X \rightarrow Y$ is a morphism of finite type then f is universally submersive iff f satisfies the weak lifting property for DVR's.

We can reformulate the definition (2.7.2) as follows:

3.8 Definition. Let X be a G -space. $f : X \rightarrow Z$ is a *topological quotient* iff f is a weak topological quotient and f satisfies the equivalent conditions of (3.7).

If Z is normal, this notion does not give anything new:

3.9 Proposition. *Let $f : X \rightarrow Z$ be a weak topological quotient. Assume that Z is normal, irreducible and every irreducible component of X dominates Z . Then $f : X \rightarrow Z$ is a topological quotient.*

Proof. We want to use the Chevalley criterion [EGA, IV.14.4.4] to show that f is universally open. We need to prove that the fiber dimension $x \mapsto \dim_x f^{-1}(f(x))$ is locally constant on X . Let $g : X \rightarrow S$ be the morphism giving the S -space structure. For $s \in S$ let G_s denote the fiber of G over s .

$x \mapsto \dim_x f^{-1}(f(x))$ and $x \mapsto \dim \text{Stab}(x)$ are always upper semi continuous. Furthermore,

$$\dim_x f^{-1}(f(x)) + \dim \text{Stab}(x) = \dim G_{g(x)}$$

is locally constant since $G \rightarrow S$ is open. Thus $x \mapsto \dim_x f^{-1}(f(x))$ is locally constant on X . $\square$

We introduce a further notion, whose main point is to allow quotients $f : X \rightarrow Z$ where $f(K) : X(K)/G(K) \rightarrow Z(K)$ is only finite-to-one. For such morphisms the conditions in (3.7) are no longer equivalent. We retain the strongest one of them.

3.10 Definition. Let X be a G -space. $f : X \rightarrow Z$ is called an *approximate quotient* if the following conditions are satisfied:

- (3.10.1) f is a G -morphism (with the trivial G -action on Z).
- (3.10.2) f is of finite type.
- (3.10.3) If K is an algebraically closed field then $f(K) : X(K)/G(K) \rightarrow Z(K)$ is surjective and finite-to-one (as a map of sets).
- (3.10.4) f is complete mod G .

Approximate quotients are stable under many operations:

3.11 Proposition. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be G -morphisms.

(3.11.1) Assume that g is finite, surjective and the G -action on Y is trivial. Then f is an approximate quotient iff $g \circ f$ is.

(3.11.2) Assume that f is finite and surjective. Then g is an approximate quotient iff $g \circ f$ is.

(3.11.3) Assume that f is an approximate (resp. topological) quotient and let $h : Y' \rightarrow Y$ be a morphism. Then $h \times f : Y' \times_Y X \rightarrow Y'$ is an approximate (resp. topological) quotient.

(3.11.4) Let $\phi : H \rightarrow G$ be a homomorphism of algebraic groups which is finite and maps onto a finite index subgroup of G . ϕ induces an H -action on X . In this case, if f is an approximate quotient by G then f is an approximate quotient by H . The converse also holds if the G -action on Y is trivial.

Proof. The last assertion is clear. In the remaining three cases the conditions (3.10.1–3) are easy to check. In the first two cases (3.6) allows us to check (3.10.4), while this is clear for (3.11.3). $\square$

The following is the main theorem of the section. It shows that approximate quotients have surprisingly good properties. The result can be viewed as a generalization of [Fogarty83]. In the complex analytic case similar questions are considered in [Roberts86].

3.12 Theorem. Let G be an affine algebraic group scheme, universally open and locally of finite type over S . Let $m : G \times X \rightarrow X$ be a proper G -action on an algebraic space X . Let $f : X \rightarrow Z$ be an approximate quotient. Then:

- (3.12.1) f is affine.
- (3.12.2) If X is of finite type over S then so is Z .
- (3.12.3) Assume in addition that G is flat over S . If F is any coherent G -sheaf on X then $(f_*F)^G$ is a coherent sheaf on Z .

3.12.4 Remark. [Fogarty83] asserts a version of this assuming only that f is submersive. Example (3.3) is a strict quotient in his sense where his assertion about coherence fails. His assertion and proof remain valid for f universally submersive.

The following consequence of (3.12) shows that topological quotients are very close to geometric quotients:

3.13 Theorem. *Let G be an affine algebraic group scheme, flat and locally of finite type over S . Let $m : G \times X \rightarrow X$ be a proper G -action on an algebraic space X . Let $f : X \rightarrow Z$ be a topological quotient. Set*

$$X/G := \operatorname{Spec}_Z(f_*\mathcal{O}_X)^G, \quad \text{and let} \quad f : X \xrightarrow{g} X/G \xrightarrow{q} Z$$

be the natural morphisms. Then $g : X \rightarrow X/G$ is the geometric quotient of X mod G and $q : X/G \rightarrow Z$ is a finite and universal homeomorphism (5.1).

Proof. By (3.12.3) $(f_*\mathcal{O}_X)^G$ is a coherent sheaf of algebras on Z , thus $q : X/G \rightarrow Z$ is finite and surjective.

$g : X \rightarrow X/G$ is a G -morphism and the G -action on X/G is trivial. g is locally of finite type since f is. If K is an algebraically closed field, the composite

$$X(K)/G(K) \rightarrow (X/G)(K) \rightarrow Z(K)$$

is an isomorphism and $(X/G)(K) \rightarrow Z(K)$ is surjective. Thus $X(K)/G(K) \rightarrow (X/G)(K)$ is an isomorphism. By (3.11.1) g is an approximate quotient, hence g is universally submersive.

Finally, $(g_*\mathcal{O}_X)^G = \mathcal{O}_{X/G}$ by construction. Thus $g : X \rightarrow X/G$ is the geometric quotient of X mod G . $\square$

3.13.1 Remark. In (3.13) assume that f is only an approximate quotient. Then g satisfies all the properties of geometric quotients except possibly (2.7.2.3). I hope to study this problem later.

3.14 Proof of (3.12). First we show that the properties (3.12.1–3) descend through finite and dominant morphisms.

The properties (3.12.2) and (3.12.1) can be studied for any morphism of schemes.

3.14.1 Lemma. [Atiyah-MacDonald69, 7.8; EGA71, 0.6.4.9] *Let $q : U \rightarrow V$ be a finite and surjective morphism of algebraic spaces. Then*

(3.14.1.1) U is (locally) of finite type over S iff V is.

(3.14.1.2) U is Noetherian iff V is. $\square$

The following is the relative version of the Chevalley theorem:

3.14.2 Lemma. [EGA71, I.7.1.4; Knutson71, p.169] *Consider a commutative diagram of algebraic spaces*

$$\begin{array}{ccc} U' & \xrightarrow{q_U} & U \\ f' \downarrow & & \downarrow f \\ V' & \xrightarrow{q_V} & V, \end{array}$$

where q_U and q_V are finite and surjective. Then f is affine iff f' is. $\square$

The descent of coherence requires a little care:

3.14.3 Lemma. *Let G be an affine algebraic group scheme, flat and of finite type over S . Let $m : G \times U \rightarrow U$ resp. $m' : G \times U' \rightarrow U'$ be proper actions with approximate quotients $f : U \rightarrow V$ resp. $f' : U' \rightarrow V'$. Assume that we have a commutative diagram*

$$\begin{array}{ccc} U' & \xrightarrow{q_U} & U \\ f' \downarrow & & \downarrow f \\ V' & \xrightarrow{q_V} & V \end{array}$$

where q_U and q_V are finite and surjective and q_U is a G -morphism. Then the following are equivalent:

(3.14.3.1) If F is any coherent G -sheaf on U then $(f_*F)^G$ is a coherent sheaf on V .

(3.14.3.2) If F' is any coherent G -sheaf on U' then $(f'_*F')^G$ is a coherent sheaf on V' .

Proof. If F' is a coherent G -sheaf on U' then $(q_U)_*F'$ is a coherent G -sheaf on U and

$$(q_V)_* ((f'_*F')^G) = (f_*((q_U)_*F'))^G.$$

This shows that (3.14.3.1) implies (3.14.3.2).

The converse is proved by induction (with respect to lexicographic ordering) on the pair $(\dim \text{Supp } F, q(F))$ where $q(F) := \sum_i \text{length}_{x_i} F$ where the x_i are the $\dim \text{Supp } F$ -dimensional generic points of $\text{Supp } F$.

We need the following result:

3.14.3.3 Claim. [SGA3, I.5.3; Seshadri77, p.264] Assume that G is flat over S . Let $0 \rightarrow F_1 \rightarrow F_2 \rightarrow F_3$ be an exact sequence of quasicoherent G -sheaves on an algebraic space with trivial G -action. Then the sequence

$$0 \rightarrow F_1^G \rightarrow F_2^G \rightarrow F_3^G$$

is exact. $\square$

Let $\mathfrak{S}$ be the class of coherent G -sheaves F such that $(f_*F)^G$ is coherent. We need to prove that $\mathfrak{S}$ is the class of all coherent G -sheaves.

3.14.3.4 Claim. Let $0 \rightarrow F_1 \rightarrow F_2 \rightarrow F_3$ be an exact sequence of coherent G -sheaves on U . If $F_1, F_3 \in \mathfrak{S}$ then $F_2 \in \mathfrak{S}$.

Proof. By (3.14.3.3) we have an exact sequence

$$0 \rightarrow (f_*F_1)^G \rightarrow (f_*F_2)^G \xrightarrow{v} (f_*F_3)^G.$$

Set $E = \text{im } v$. If $(f_*F_3)^G$ is coherent then $E \subset (f_*F_3)^G$ is also coherent. If in addition $(f_*F_1)^G$ is coherent then $(f_*F_2)^G$ is also coherent. $\square$

Let F be a coherent G -sheaf on U . By induction, if E is a coherent G -sheaf on U such that $(\dim \text{Supp } E, q(E)) < (\dim \text{Supp } F, q(F))$ then $E \in \mathfrak{S}$. We need to prove that $F \in \mathfrak{S}$.

Let E denote the kernel of the natural map $p : F \rightarrow (q_U)_*(q_U^*F)$. We have an exact sequence

$$0 \rightarrow E \rightarrow F \xrightarrow{p} (q_U)_*(q_U^*F).$$

p is nonzero at all generic points of $\text{Supp } F$, thus $(\dim \text{Supp } E, q(E)) < (\dim \text{Supp } F, q(F))$, hence $E \in \mathfrak{S}$. Set $F' := q_U^*F$. Then

$$(q_V)_* ((f'_*F')^G) = (f_*((q_U)_*F'))^G = (f_*((q_U)_*(q_U^*F)))^G.$$

By (3.14.3.1) $(f'_*F')^G$ is coherent on V' , thus $(q_V)_* ((f'_*F')^G)$ is coherent. This shows that $(q_U)_*(q_U^*F) \in \mathfrak{S}$. By (3.14.3.4) this implies that $(f_*F)^G$ is coherent. $\square$

The rest of the proof of (3.12) relies on a slight strengthening of a result of Mumford (see [EGA, IV, 14.5.10]).

3.14.4 Proposition. *Let $f : X \rightarrow Z$ be a universally open and surjective morphism, locally of finite type. Then there is a finite and surjective morphism $g : Z' \rightarrow Z$ and an open cover $\cup Z'_i = Z'$ such that for every irreducible component $X'_{ij} \subset X'_i := X \times_Z Z'_i$ the induced morphism*

$$f'_{ij} : X'_{ij} \rightarrow Z'_i \quad \text{has a section} \quad s_{ij} : Z'_i \rightarrow X'_{ij} \quad \text{for every } i, j. \quad \square$$

Apply (3.14.4) to $f : X \rightarrow Z$ in (3.12) and set $X' := X \times_Z Z'$. $f' : X' \rightarrow Z'$ is an approximate quotient by (3.11.3). By (3.14.1–3) it is sufficient to prove (3.12) for $f' : X' \rightarrow Z'$.

The conditions (3.12.1–3) are local on Z' , thus we may replace Z' by Z'_i to obtain $f'_i : X'_i \rightarrow Z'_i$. Let $X'_{ij} \subset X'_i$, $j = 1, \dots, j(i)$ be the irreducible components. For every i, j we get a morphism

$$p_{ij} : G \times_S Z'_i \xrightarrow{(id_G, s_{ij})} G \times_S X'_i \xrightarrow{m} X'_{ij}.$$

Let Z''_i denote the disjoint union of $j(i)$ copies of Z'_i . The sum of the morphisms p_{ij} gives a finite and surjective G -morphism $p_i := \sum_j p_{ij}$ and a commutative diagram

$$\begin{array}{ccc} G \times_S Z''_i & \xrightarrow{p_i} & X'_i \\ \downarrow & & \downarrow f'_i \\ Z''_i & \longrightarrow & Z'_i. \end{array}$$

(3.14.1–3) show that it is sufficient to prove the following:

3.14.5 Lemma. *(3.12) is true in case $X = G \times_S Z$ and $f : X \rightarrow Z$ is the second projection.*

Proof. Since G/S is affine, so is $G \times_S Z \rightarrow Z$. If $e : S \rightarrow G$ denotes the unit section then $Z \cong S \times_S Z \hookrightarrow X$, so Z is of finite type if X is. These show (3.12.1) and (3.12.2).

$f : X \rightarrow Z$ is a group scheme over Z . Thus $E \mapsto f^*E$ provides an equivalence of the category of (quasi)coherent sheaves on Z and the category of (quasi)coherent G -sheaves on X . Its inverse is given by $f^*E \mapsto (f_*(f^*E))^G = E$. Thus if F is a coherent G -sheaf on X then $(f_*F)^G$ is a coherent sheaf on Z . $\square$

This completes the proof of (3.12). $\square$

4. QUOTIENTS OF NORMAL SPACES BY SESHADRI

The aim of this section is to consider quotients of normal algebraic spaces by proper actions of algebraic group schemes. This situation has been studied in [Seshadri63,72] and his methods give an almost complete answer.

We need to apply his techniques in a more general setting. Therefore we give precise statements. The proofs require only slight technical modifications. Some of these are worked out in [Kollár95a].

4.1 Proposition. [Seshadri72, Lemma 6.1] *Let G be an algebraic group scheme, smooth and of finite type over S . Let X be an algebraic G -space and $n : \bar{X} \rightarrow X$ the normalisation. Assume that n is finite.*

Then $\bar{X}$ has a unique structure of an algebraic G -space such that n is a G -morphism. $\square$

4.1.1 Remark. There are some natural group schemes over $\mathbb{Z}$ which are not smooth over $\mathbb{Z}$. The simplest example is the group of square roots of unity $\mu_2 := \text{Spec } \mathbb{Z}[x]/(x^2 - 1)$. This has two irreducible components (corresponding to $+1$ and -1) which intersect in characteristic 2. Its normalization consists of two copies of $\text{Spec } \mathbb{Z}$, and the action of μ_2 on itself does not lift to an action on the normalization.

4.2 Theorem. [Seshadri72, Theorem 6.1] *Let X be an excellent algebraic space over S and G an affine algebraic group scheme, smooth and of finite type over S . Let $m_X : G \times X \rightarrow X$ be a proper G -action on X .*

Then there is a Zariski locally trivial principal G -bundle $p : Y \rightarrow Z(Y)$ and a finite and dominant G -morphism $q : Y \rightarrow X$.

If X is irreducible, then we can choose $p : Y \rightarrow Z(Y)$ and $q : Y \rightarrow X$ in such a way that Y is irreducible, the field extension $k(Y) \supset k(X)$ is normal with Galois group Γ and the action of Γ on Y commutes with the G -action on Y . $\square$

As a consequence we obtain the first quotient theorem:

4.3 Theorem. *Let G be an affine algebraic group scheme, smooth and of finite type over S and X a normal algebraic space of finite type over S . Let $m : G \times X \rightarrow X$ be a proper G -action on X .*

Then the geometric quotient $f : X \rightarrow X/G$ exists, and X/G is a normal algebraic space of finite type over S .

Proof. We may assume that X is irreducible. Choose $q : Y \rightarrow X$ and $p : Y \rightarrow Z(Y)$ as in (4.2). Set $Y' = Y/\Gamma$ and $Z' = Z(Y)/\Gamma$; these exist by (4.4). p descends to a morphism $p' : Y' \rightarrow Z'$. Since the G and the Γ actions commute, the G -action on Y descends to a G -action on Y' . From (3.11) we conclude that p' is a topological quotient of Y' mod G .

Assume next that $k(X)$ has characteristic zero. Then $k(X) = k(Y)^\Gamma$, hence $Y' = Y$ and $Z = Z'$ is a topological quotient of Y mod G . Thus a geometric quotient exists by (3.13).

If $k(X)$ has positive characteristic then $Y' \rightarrow Y$ is a purely inseparable G -morphism. A topological quotient of Y' mod G exists. By (6.7) this implies that a topological quotient of Y mod G also exists. $\square$

In the proof we used that quotients of algebraic spaces by finite groups exist. This result is due to Deligne (cf. [Knutson71, p.185]), but I do not know of any published proof. In the case that is used here, this also follows from (4.7).

4.4 Lemma. *Let X be a separated algebraic space of finite type over S and G a finite group acting on X . Then a geometric quotient X/G exists and X/G is also of finite type over S . $\square$*

It is not more difficult to understand quotients of normal algebraic spaces by finite equivalence relations. Their existence is established through the following two constructions.

4.5 Construction of Symmetric Products. Let X be a separated algebraic space. Our aim is to construct its symmetric products, denoted by $S^n X$. X can be given by an étale equivalence relation $R \rightrightarrows U$ where U is affine. Since X is separated, the

resulting morphism $\phi : R \rightarrow U \times U$ is a closed immersion. We can write $R = \Delta \cup R'$ (disjoint union) where $\phi(\Delta)$ is the diagonal.

The induced morphism $f^n : U^n \rightarrow X^n$ is étale. The symmetric group S_n acts on both spaces and f^n is S_n -equivariant. Unfortunately, it is not fixed-point reflecting. More precisely, it is not fixed-point reflecting at $(x_1, \dots, x_n)$ iff there is a pair $i < j$ such that $f(x_i) = f(x_j)$ and $x_i \neq x_j$. Set

$$W_n := U^n \setminus \bigcup_{i < j} \pi_{ij}^{-1}(\phi(R')),$$

where $\pi_{ij} : U^n \rightarrow U^2$ is the projection to the product of the i^{th} and j^{th} factors. $f^n : W_n \rightarrow X^n$ is étale, surjective and fixed-point reflecting. W_n is quasi affine, hence the geometric quotient W_n/S_n exists and is also quasi affine (cf. [Mumford68, Sec. 12]). Thus $S^n X$ is given by the equivalence relation

$$(W_n \times_{X^n} W_n)/S_n \rightrightarrows W_n/S_n.$$

4.6 Lemma. *Let X, Y be separated algebraic spaces, X normal and irreducible. Let $Z \subset X \times Y$ be a closed subspace such that the projection $p_X : Z \rightarrow X$ is finite and surjective on every irreducible component of Z . Let $d = \deg(Z/X)$. Then there is a natural morphism $\phi_Z : X \rightarrow S^d Y$.*

Proof. Let $X^0 \subset X$ be a dense open set such that $p_X|Z$ is flat over X^0 . Then

$$x \mapsto p_Y[p_X^{-1}(x)] \in S^d Y$$

is a well defined morphism. By looking at the closure of the graph it is easy to see that it extends to $\phi_Z : X \rightarrow S^d Y$ such that $\text{Supp } \phi_Z(x) = \text{Supp } p_Y[p_X^{-1}(x)]$ for every $x \in X$. (In general no such extension exists if X is not normal.) $\square$

4.7 Corollary. *Let X be a normal algebraic space. Let $Z \subset X \times X$ be a closed subspace such that the first projection $p_X : Z \rightarrow X$ is finite and surjective on every irreducible component of Z . Assume that Z defines an equivalence relation on X . (For the present purposes all we need is that it is an equivalence relation on closed points in algebraically closed fields.) Then there is a morphism $\phi_Z : X \rightarrow X/Z$ whose fibers are precisely the Z -equivalence classes.*

Proof. Set $d = \deg(Z/X)$ and let X/Z be the normalization of the image of $\phi_Z : X \rightarrow S^d X$. $\square$

4.7.1 Remark. The above construction gives the correct fiber set theoretically, but not necessarily scheme theoretically, especially in positive characteristic. Thus (4.7) implies that if X is a normal algebraic space and G a finite and flat group scheme acting on X then a topological quotient exists. Thus by (3.13) a geometric quotient also exists.

5. WEAKLY NORMAL SPACES

The aim of this section is to consider the quotient problem for weakly normal algebraic spaces. The notion of weak normality and of universal homeomorphisms are central to the construction of quotients. We start by recalling their definitions and basic properties.

5.1 Definition. We say that a morphism of schemes $g : U \rightarrow V$ is a *universal homeomorphism* if it is a homeomorphism and for every $W \rightarrow V$ the induced morphism $U \times_V W \rightarrow W$ is again a homeomorphism.

This notion extends to morphisms of algebraic spaces the usual way [Knutson71, II.3].

This concept is especially useful if g is finite. In this case we have the following characterization (cf. [EGA71, I.3.7–8]).

5.2 Lemma. *Let $g : U \rightarrow V$ be a finite and surjective morphism of algebraic spaces. The following are equivalent*

(5.2.1) *g is a universal homeomorphism.*

(5.2.2) *g is surjective and universally injective.*

(5.2.3) *For every $v \in V$ the fiber $g^{-1}(v)$ has a single point v' and $k(v')$ is a purely inseparable field extension of $k(v)$. (Thus $k(v') = k(v)$ if $\text{char } k(v) = 0$.)* $\square$

5.2.4 Remark. The terminology in English does not seem to be uniform. The french “radiciel” [EGA71, I.3.7] did not catch on. Some authors use the name “purely inseparable morphism” to refer to a morphism satisfying the equivalent properties in (5.2).

One of the most important properties of these morphisms is the following:

5.3 Theorem. [SGA1, IX.4.10] *Let $g : U \rightarrow V$ be a finite and universal homeomorphism. Then $* \mapsto * \times_V U$ induces an equivalence between the categories*

$$(\text{étale morphisms: } * \rightarrow V) \xrightarrow{* \mapsto * \times_V U} (\text{étale morphisms: } * \rightarrow U). \quad \square$$

The notion of weak normality was developed in a series of papers [Andreotti-Noguet67; Andreotti-Bombieri69; Traverso70]. We recall the definitions and a few of the basic properties. A short outline of the theory can also be found in [Kollár95b, I.7].

5.4 Definition. The *weak normalization* of an algebraic space X is an algebraic space X' together with a finite and surjective morphism $m : X' \rightarrow X$ such that

(5.4.1) for every $x \in X$ there is a unique $x' \in X'$ such that $m(x') = x$;

(5.4.2) for every $x \in X$ the field extension $k(x') \supset k(x)$ is purely inseparable and an isomorphism if $x \in X$ is a generic point;

(5.4.3) if $g : Y \rightarrow X$ satisfies the above two properties then there is a unique factorization

$$m : X' \rightarrow Y \xrightarrow{g} X.$$

An algebraic space X is called *weakly normal* if it is its own weak normalization.

5.5 Proposition. [Andreotti-Noguet67, Andreotti-Bombieri69, Traverso70] *Let X be an algebraic space such that its normalization $\bar{X} \rightarrow X$ is finite over X . Then:*

(5.5.1) *X has a weak normalization $n : X^{\text{wn}} \rightarrow X$. n is a finite and universal homeomorphism.*

(5.5.2) *There is a factorization $\bar{X} \rightarrow X^{\text{wn}} \rightarrow X$.*

(5.5.3) *The conductor ideal (5.7) $I_{\bar{X}/X^{\text{wn}}} \subset \mathcal{O}_{\bar{X}}$ is its own radical.*

(5.5.4) *Let $f : Y \rightarrow X$ be a morphism such that X is weakly normal at the image of every generic point of Y . Then f lifts to a morphism $f^{\text{wn}} : Y^{\text{wn}} \rightarrow X^{\text{wn}}$.*

(5.5.5) *If $f : Y \rightarrow X$ is smooth and X is weakly normal then so is Y .*

Proof. For the reader's convenience, let us recall the construction of X^{wn} for schemes.

For a point $x \in X$ let $\bar{x} = n^{-1}(x) \subset \bar{X}$ with reduced scheme structure. $k(\bar{x})$ is the direct sum of the residue fields of the points of $\bar{x}$, $i_x : k(x) \rightarrow k(\bar{x})$ the natural injection. For an open affine $U \subset X$ let

$$\mathcal{O}^{\text{wn}}(U) := \{f \in \mathcal{O}_{n^{-1}(U)} \mid f(\bar{x})^q \in \text{im } i_x \ \forall x \in U, \text{ where } q \text{ is a power of } \text{char } k(x).\}$$

This defines a sheaf of rings $\mathcal{O}^{\text{wn}}$ over X . Set $X^{\text{wn}} = \text{Spec}_X \mathcal{O}^{\text{wn}}$.

The properties (5.5.1–4) for schemes are clear from this construction. (5.5.5) is in [Kollár95b, I.7.2.6].

Let $u : U \rightarrow V$ be a surjective étale morphism of schemes. By (5.5.5) U is weakly normal iff V is. Thus all the above assertions descend from the category of schemes to the category of algebraic spaces (cf. [Knutson71, p.106]). $\square$

5.6 Corollary. *Let G be an algebraic group scheme, smooth and of finite type over S . Let X be an algebraic G -space and $n : X^{\text{wn}} \rightarrow X$ the weak normalization. Assume that n is finite.*

Then X^{wn} has a unique structure of an algebraic G -space such that n is a G -morphism.

Proof. The G -action on X is given by a morphism $m : G \times X \rightarrow X$. Since G is smooth over S , (5.5.5) implies that $G \times (X^{\text{wn}}) = (G \times X)^{\text{wn}}$. Thus by (5.5.4) m lifts to a morphism

$$m^{\text{wn}} : G \times X^{\text{wn}} \rightarrow X^{\text{wn}}.$$

It is clear that m^{wn} defines a group action. $\square$

Our aim is to construct geometric quotients of weakly normal G -spaces by comparing the situation to the normalization. Let $\bar{X} \rightarrow X$ be the normalization. Assume that the G -action lifts to an action on $\bar{X}$ and that a geometric quotient $\bar{X}/G$ exists. Thus we have a diagram

$$\begin{array}{ccc} \bar{X} & \longrightarrow & X \\ \downarrow & & \\ \bar{X}/G & & \end{array}$$

and X/G would complete it to a commutative square. The construction of X/G becomes a special case of constructing push-outs (8.1).

Before stating the main result of this section, we need to introduce further notation:

5.7 Notation. Let G/S be an algebraic group scheme, flat over S . Let X/S be a weakly normal algebraic space over S with normalization $p : \bar{X} \rightarrow X$ which is assumed to be finite over X . Assume that we are given G -actions on $\bar{X}$ and on X such that n is a G -morphism.

Let $I_X \subset \mathcal{O}_X$ resp. $I_{\bar{X}} \subset \mathcal{O}_{\bar{X}}$ denote the *conductor ideals*. That is,

This definition shows that I_X and $I_{\bar{X}}$ are G -sheaves. The corresponding subschemes are denoted by $C_X \subset X$ resp. $C_{\bar{X}} \subset \bar{X}$. By (5.5.3) I_X and $I_{\bar{X}}$ are their own radicals, so the conductor subschemes are reduced. The G action restricts to G -actions on C_X resp. $C_{\bar{X}}$. p restricts to a finite and surjective G -morphism $p : C_{\bar{X}} \rightarrow C_X$.

Assume that a geometric quotient $\bar{f} : \bar{X} \rightarrow Z(\bar{X})$ exists. $C_{\bar{X}} \subset \bar{X}$ is G -invariant and closed, thus there is a closed and reduced subscheme $j : Z(\bar{C}) \hookrightarrow Z(\bar{X})$ such that $\bar{f}^{-1}(Z(\bar{C})) = C_{\bar{X}}$ (as sets, but not necessarily as schemes).

By (3.11.3) $C_{\bar{X}} \rightarrow Z(\bar{C})$ is a topological quotient of $C_{\bar{X}}$ by G . Therefore, by (3.13) a geometric quotient $C_{\bar{X}} \rightarrow Z(C_{\bar{X}})$ exists and the induced morphism $q_1 : Z(C_{\bar{X}}) \rightarrow Z(\bar{C})$ is a finite and universal homeomorphism.

Assume furthermore that a geometric quotient $C_X \rightarrow Z(C_X)$ also exists. Since $p : C_{\bar{X}} \rightarrow C_X$ is a G -morphism, we obtain an induced morphism $q_2 : Z(C_{\bar{X}}) \rightarrow Z(C_X)$.

5.8 Theorem. *Let G be an algebraic group scheme, flat and of finite type over S . Let X be a weakly normal algebraic G -space of finite type over S such that the G -action lifts to a G -action on the normalization $p : \bar{X} \rightarrow X$. Assume that:*

(5.8.1) *The geometric quotients $\bar{f} : \bar{X} \rightarrow Z(\bar{X})$ and $C_X \rightarrow Z(C_X)$ exist.*

(5.8.2) *The push-out diagram*

$$(5.8.2.1) \quad \begin{array}{ccc} Z(C_{\bar{X}}) & \xrightarrow{q_1} & Z(\bar{C}) \\ q_2 \downarrow & & \\ Z(C_X) & & \end{array}$$

can be completed to

$$(5.8.2.2) \quad \begin{array}{ccc} Z(C_{\bar{X}}) & \xrightarrow{q_1} & Z(\bar{C}) \\ q_2 \downarrow & & \downarrow s_2 \\ Z(C_X) & \xrightarrow{s_1} & W, \end{array}$$

such that s_2 is finite and s_1 is a finite and universal homeomorphism.

Then a geometric quotient of X mod G exists.

Proof. Assume that s_2 is finite. By (8.6) there is an algebraic space Z and finite morphisms $q : Z(\bar{X}) \rightarrow Z$ and $j_W : W \hookrightarrow Z$ which fit into the following commutative diagram

$$\begin{array}{ccc} Z(\bar{C}) & \xrightarrow{j} & Z(\bar{X}) \\ s_2 \downarrow & & \downarrow q \\ W & \xrightarrow{j_W} & Z. \end{array}$$

The next two claims imply (5.8):

5.8.3 Claim. (5.8.3.1) $\bar{f} : \bar{X} \rightarrow Z(\bar{X})$ descends to a G -morphism $f : X \rightarrow Z$.

(5.8.3.2) $f : X \rightarrow Z$ is a topological quotient of X by G .

Proof. The first assertion means that the diagram

$$\begin{array}{ccc} \bar{X} & \xrightarrow{p} & X \\ \bar{f} \downarrow & & \\ Z(\bar{X}) & \xrightarrow{q} & Z \end{array}$$

can be completed to a push-out diagram by $f : X \rightarrow Z$. At the set theoretic level this defines f uniquely as $f = q \circ \bar{f} \circ p^{-1}$.

We construct f by constructing its graph. The morphism $p \times (q \circ \bar{f}) : \bar{X} \rightarrow X \times Z$ is finite; let $Y \subset X \times Z$ denote its image (with reduced scheme structure).

Y is the graph of the (a priori multi-valued) map $q \circ \bar{f} \circ p^{-1}$. If the projection $h : Y \rightarrow X$ is an isomorphism, then it defines a morphism $f : X \rightarrow Z$. X is weakly normal and Y is dominated by the normalization $\bar{X}$, thus by (5.5) it is sufficient to prove that $h : Y \rightarrow X$ is universally injective.

Given $x \in X$ let $\text{red } p^{-1}(x) = \bar{x}_1 \cup \cdots \cup \bar{x}_s$ (where s depends on x). Then

$$\text{red } h^{-1}(x) = \bigcup_i (x, (q \circ \bar{f})(\bar{x}_i)).$$

If $x \notin C_X$ then $\bar{X} \rightarrow X$ is an isomorphism over x , thus $Y \rightarrow X$ is also an isomorphism over x .

Next assume that $x \in C_X$, hence $\text{red } p^{-1}(x) \subset C_{\bar{X}}$. $q \circ \bar{f} : C_{\bar{X}} \rightarrow Z$ sits in a commutative diagram

$$(5.8.3.3) \quad \begin{array}{ccccccc} C_{\bar{X}} & \xrightarrow{\bar{f}} & Z(C_{\bar{X}}) & \xrightarrow{q_1} & Z(\bar{C}) & \xrightarrow{j} & Z(\bar{X}) \\ p \downarrow & & q_2 \downarrow & & \downarrow s_2 & & q \downarrow \\ C_X & \longrightarrow & Z(C_X) & \xrightarrow{s_1} & W & \xrightarrow{j_W} & Z. \end{array}$$

This shows that $(q \circ \bar{f})(\bar{x}_i)$ is independent of i and the residue field of Z at $(q \circ \bar{f})(\bar{x}_i)$ is a subfield of the residue field of X at x . Therefore $h : Y \rightarrow X$ is universally injective, hence an isomorphism. This shows (5.8.3.1).

The morphisms $p : \bar{X} \rightarrow X$ and $q : Z(\bar{X}) \rightarrow Z$ are finite, thus Z is an approximate quotient of $X \bmod G$ by (3.11).

In order to see that it is a topological quotient we need to check that for every algebraically closed field K (2.7.2.3) holds. By construction

$$\begin{aligned} X(K)/G(K) &= (X - C_X)(K)/G(K) \cup C_X(K)/G(K) \\ &= (\bar{X} - C_{\bar{X}})(K)/G(K) \cup C_X(K)/G(K) \\ &= (Z(\bar{X}) - Z(\bar{C}))(K) \cup Z(C_X)(K) \\ &= (Z - W)(K) \cup W(K) = Z(K). \end{aligned}$$

All the unions are disjoint, thus $X(K)/G(K) \rightarrow Z(K)$ is an isomorphism of sets. $\square$

6. QUOTIENTS IN POSITIVE CHARACTERISTIC

In this section, k denotes a field of positive characteristic p . The main result of the section is the following:

6.1 Theorem. *Let G be an affine algebraic group scheme of finite type over k and X an algebraic space of finite type over k . Let $m : G \times X \rightarrow X$ be a proper G -action on X .*

Then a geometric quotient $f : X \rightarrow X/G$ exists, and X/G is an algebraic space of finite type over k .

We already proved this result in case X is normal (4.2). In this section we reduce the general case to the normal case.

The crucial feature of positive characteristic algebraic spaces that we use is the Frobenius morphism:

6.2 Definition of the Geometric Frobenius Morphism [SGA5, XIV].

Let S be an $\mathbb{F}_p$ -scheme. Fix $q = p^r$ for some natural number r . Then $a \mapsto a^q$ is an endomorphism of $\mathcal{O}_S$. This can be extended to polynomials by the formula

$$f = \sum a_I x^I \mapsto f^{(q)} := \sum a_I^q x^I.$$

Let $U = \text{Spec } R$ be an affine scheme over S . Write $R = \mathcal{O}_S[x_1, \dots, x_m]/(f_1, \dots, f_n)$ and set

$$R^{(q)} := \mathcal{O}_S[x_1^{(q)}, \dots, x_m^{(q)}]/(f_1^{(q)}, \dots, f_n^{(q)}) \quad \text{and} \quad U^{(q)} := \text{Spec } R^{(q)},$$

where the $x_i^{(q)}$ are new variables. There are natural morphisms

$$F^q : U \rightarrow U^{(q)} \quad \text{and} \quad (F^q)^* : R^{(q)} \rightarrow R \quad \text{given by} \quad (F^q)^*(x_i^{(q)}) = x_i^q.$$

It is easy to see that these are independent of the choices made. Thus F^q gives a functor from algebraic spaces over S to algebraic spaces over S .

If $F^q : \text{Spec } S \rightarrow \text{Spec } S$ denotes the q^{th} -power morphism, then one can define $X^{(q)}$ intrinsically as

$$X^{(q)} = X \times_{\text{Spec } S, F^q} \text{Spec } S.$$

Since $X^{(q)}$ is obtained by base change, it commutes with many other constructions. In particular, we obtain the following:

6.3 Lemma. *Let $m : G \times X \rightarrow X$ be a G -action.*

(6.3.1) *$m^{(q)} : G^{(q)} \times X^{(q)} \rightarrow X^{(q)}$ defines a $G^{(q)}$ -action on $X^{(q)}$.*

(6.3.2) *m is proper iff $m^{(q)}$ is.*

(6.3.3) *$f : X \rightarrow Z$ is a topological quotient of X by G iff $f^{(q)} : X^{(q)} \rightarrow Z^{(q)}$ is a topological quotient of $X^{(q)}$ by $G^{(q)}$. $\square$*

Frobenius morphisms give the most interesting examples of universal homeomorphisms:

6.4 Lemma. *Let X be an algebraic space of finite type over S . Then $F^q : X \rightarrow X^{(q)}$ is a finite and universal homeomorphism. $\square$*

6.5 Remark. There are many algebraic spaces which are not of finite type and such that $F^q : X \rightarrow X^{(q)}$ is finite. By a result of Kunz (see [Matsumura80, p.302]) such algebraic spaces are excellent. The results of this section probably remain valid for algebraic spaces satisfying this property. The following example of [Nagata69] shows that this assumption is necessary.

6.5.1 Example. Let k be a field of characteristic $p > 0$ and $K := k(x_1, x_2, \dots)$, where the x_i are algebraically independent over k . Let

$$D := \sum x_i^{1+p} \frac{\partial}{\partial x_i} \quad \text{be a derivation of } K.$$

Let $F := \{f \in K \mid D(f) = 0\}$ be the subfield of constants. Set

$$R = K + \epsilon K \quad \text{where } \epsilon^2 = 0, \text{ and } \sigma : f + \epsilon g \mapsto f + \epsilon(g + D(f)).$$

R is a local Artin ring. It is easy to check that σ is an automorphism of R of order p . The fixed ring is $R^\sigma = F + \epsilon K$. Its maximal ideal is $m := (\epsilon K)$ and generating sets of m correspond to bases of K as an F -vectorspace. (6.5.2) shows that m is not finitely generated.

6.5.2 Claim. *Notation as above. The x_i are linearly independent over F .*

Proof. Assume that we have a relation $\sum_{i \in I} f_i x_i = 0$ where $|I|$ is the smallest possible. Apply D to get that

$$0 = \sum_{i \in I} f_i D(x_i) = \sum_{i \in I} (x_i^p f_i) x_i.$$

$x_i^p \in F$, thus we get another linear dependence relation, a contradiction. $\square$

For us the most important feature of the Frobenius morphism is the following universal property:

6.6 Proposition. *Let X, Y be algebraic spaces over S and $g : X \rightarrow Y$ a finite and universal homeomorphism. Then there is a $q = p^r$ such that F^q can be factored as*

$$F^q : X \xrightarrow{g} Y \xrightarrow{\bar{g}} X^{(q)}.$$

For large q the factorization is canonical in the sense that if

$$\begin{array}{ccc} X_1 & \xrightarrow{g_1} & Y_1 \\ f \downarrow & & h \downarrow \\ X_2 & \xrightarrow{g_2} & Y_2 \end{array}$$

is a commutative diagram where the g_i are finite and universal homeomorphisms, then the factorization gives a commutative diagram

$$\begin{array}{ccccc} X_1 & \xrightarrow{g_1} & Y_1 & \xrightarrow{\bar{g}_1} & X_1^{(q)} \\ f \downarrow & & h \downarrow & & f^{(q)} \downarrow \\ X_2 & \xrightarrow{g_2} & Y_2 & \xrightarrow{\bar{g}_2} & X_2^{(q)}. \end{array}$$

Proof. Since the factorization is canonical, it is sufficient to construct it in case X and Y are affine schemes over an affine scheme $\text{Spec } C$.

The latter can be formulated in terms of a ring homomorphism $g^* : A \rightarrow B$ where A and B are C -algebras. We can decompose g^* into $A \twoheadrightarrow B_1$ and $B_1 \hookrightarrow B$. We deal with them separately.

First consider $B_1 \subset B$. We need to show that there is a q such that $B^q \subset B_1$, where B^q denotes the C -algebra generated by the q -th powers of all elements. The proof is by Noetherian induction.

If B_1 is Artin, then the assertion is clear. Applying this over the generic points we obtain that $B_1 \subset B_1 B^q$ is an isomorphism at all generic points for $q \gg 1$. Let $I < B_1$ denote the conductor of this extension.

By induction we know that there is a q' such that $(B_1 B^q / I)^{q'} \subset B_1 / I$. Thus we get

$$B^{(qq')} \rightarrow B^{qq'} \subset (B_1 B^q)^{q'} \subset B_1.$$

Next consider $A \twoheadrightarrow B_1$. The kernel is a nilpotent ideal $I < A$, say $I^m = 0$. Choose q' such that $q' \geq m$. For $b_1 \in B_1$ let $b'_1 \in A$ be any preimage. Then $(b'_1)^{q'}$ depends only on b_1 . The map

$$b_1 \mapsto (b'_1)^{q'} \quad \text{defines a factorization} \quad B_1^{(q')} \rightarrow A \rightarrow B_1.$$

Combining the map $B^{(q)} \rightarrow B_1$ with $B_1^{(q')} \rightarrow A$ we obtain $B^{(qq')} \rightarrow A$. $\square$

The main technical result of the section is the following:

6.7 Theorem. *Let G, H be algebraic group schemes of finite type over S and $\rho : H \rightarrow G$ a group homomorphism which is a finite and universal homeomorphism. Let Y, X be algebraic spaces of finite type over S and $f : Y \rightarrow X$ a finite and universal homeomorphism.*

Let $m_Y : H \times Y \rightarrow Y$ and $m_X : G \times X \rightarrow X$ be proper actions and assume that f is a ρ -morphism, that is, the following diagram is commutative

$$(6.7.1) \quad \begin{array}{ccc} H \times Y & \xrightarrow{\rho \times f} & G \times X \\ m_Y \downarrow & & m_X \downarrow \\ Y & \xrightarrow{f} & X. \end{array}$$

Then the following are equivalent:

(6.7.2) *The geometric quotient of X mod G exists.*

(6.7.3) *The geometric quotient of Y mod H exists.*

Proof. Let $p_X : X \rightarrow X/G$ be the geometric quotient. Then $p_X \circ f : Y \rightarrow X/G$ is a topological quotient by (3.11), hence a geometric quotient of Y mod H exists by (3.13).

Conversely, let $p_Y : Y \rightarrow Y/H$ be the geometric quotient. By (6.3.3) $p_Y^{(q)} : Y^{(q)} \rightarrow (Y/H)^{(q)}$ is a topological quotient of $Y^{(q)}$ mod $H^{(q)}$. By (6.6) the diagram (6.7.1) can be completed to

$$(6.7.4) \quad \begin{array}{ccccc} H \times Y & \xrightarrow{\rho \times f} & G \times X & \xrightarrow{\bar{\rho} \times \bar{f}} & H^{(q)} \times Y^{(q)} \\ m_Y \downarrow & & m_X \downarrow & & m_Y^{(q)} \downarrow \\ Y & \xrightarrow{f} & X & \xrightarrow{\bar{f}} & Y^{(q)}. \end{array}$$

$p_Y^{(q)} \circ \bar{f} : X \rightarrow (Y/H)^{(q)}$ is a topological quotient as before, hence a geometric quotient of X mod G exists by (3.13). $\square$

6.8 Corollary. *Let G be an affine algebraic group scheme of finite type over k and X an algebraic G -space of finite type over k . Assume that the G -action on X is proper.*

Then, a geometric quotient of $X \bmod G$ exists iff a geometric quotient of $X \bmod \text{red } G$ exists.

Proof. Apply (6.7) with $Y = X$ and $H = \text{red } G$. $\square$

6.9 Corollary. *Let G be a smooth, affine algebraic group scheme of finite type over k and X an algebraic G -space of finite type over k . Assume that the G -action on X is proper.*

Then, a geometric quotient of $X \bmod G$ exists iff a geometric quotient of $(\text{red } X)^{\text{wn}} \bmod G$ exists.

Proof. Apply (6.7) with $Y = (\text{red } X)^{\text{wn}}$ and $H = G$. The G action on X gives a G -action on $\text{red } X$ and this in turn gives a G -action on $(\text{red } X)^{\text{wn}}$ by (5.6). (Both times we used that G is smooth.) $\square$

6.10 Corollary. *Let G be a smooth, affine algebraic group scheme of finite type over k and X a weakly normal algebraic G -space of finite type over k . Assume that the G -action on X is proper. Let $p : \bar{X} \rightarrow X$ denote the normalization.*

Then, the geometric quotient of X/G exists if $\bar{X}/G$ exists.

Proof. Assume that $\bar{f} : \bar{X} \rightarrow \bar{X}/G$ exists. We use the notation of (5.7).

$Z(C_{\bar{X}}) \rightarrow Z(\bar{C})$ is a finite and universal homeomorphism by (3.11.3). Hence by (8.4) a push-out diagram as in (5.8.2.2) exists. Thus a geometric quotient of $X \bmod G$ exists by (5.8). $\square$

6.11 Proof of (6.1). Start with arbitrary G and X . We would like to construct a geometric quotient of X by G . By (6.3.3) it is sufficient to show that the geometric quotient $X^{(q)}/G^{(q)}$ exists for some q . If $q \gg 1$ then $\text{red } G^{(q)}$ is smooth over k by [Weil62, p.25]. Thus by (6.8) it is sufficient to show the existence of a geometric quotient in case G is smooth over k .

By (6.9) this can be further reduced to the case when in addition X is weakly normal, and by (6.10) to the case when X is normal. The latter types of actions have already been dealt with in (4.2). $\square$

The methods of this section also apply to group schemes in positive characteristic. The only part that fails is the reduction to smooth group schemes in (6.11). We obtain the following result:

6.12 Theorem. *Let S be an excellent scheme of characteristic p . Let G be a smooth, affine algebraic group scheme of finite type over S and X an algebraic space of finite type over S . Let $m : G \times X \rightarrow X$ be a proper G -action on X .*

Then the geometric quotient $f : X \rightarrow X/G$ exists, and X/G is an algebraic space of finite type over S . $\square$

7. QUOTIENTS BY REDUCTIVE GROUP SCHEMES

7.1 Definition. A group scheme G/S is called reductive if G is smooth over S and for every field $\text{Spec } K \rightarrow S$ the fiber product $G_K := G \times_S \text{Spec } K$ is a connected and reductive algebraic group over K .

See [Demazure67] for a detailed study of the structure of reductive group schemes. $\blacksquare$

7.2 Examples. The simplest and for moduli problems the most important reductive group schemes are $SL(n, \mathbb{Z})$ and $PGL(n, \mathbb{Z})$, as group schemes over $\mathbb{Z}$. For a field K the corresponding algebraic groups are $SL(n, K)$ resp. $PGL(n, K)$.

[Seshadri77] studied the invariant theory of reductive group schemes. Two of his results are crucial for our purposes. (As always, we assume that S is excellent.)

7.3 Theorem. [Seshadri77] *Let G/S be a reductive group scheme. Let X/S be an affine scheme of finite type over S and $m : G \times X \rightarrow X$ a G -action. Assume for simplicity that the action is proper. Then the following hold.*

(7.3.1) *The geometric quotient $f : X \rightarrow X/G$ exists and X/G is an affine scheme of finite type over S .*

(7.3.2) *If S is a scheme over $\text{Spec } \mathbb{Q}$ then $f : X \rightarrow X/G$ is a universal geometric quotient. That is, if $g : Y \rightarrow X$ is a G -morphism which is a closed immersion then the induced morphism $f/G : Y/G \rightarrow X/G$ is also a closed immersion.*

Proof. The first assertion is [Seshadri77, p.263]. The second part is not mentioned explicitly, so I outline how it follows from the results of the article.

Pushing forward to the base scheme S , (7.3.2) is equivalent to the following assertion:

7.3.3 Claim. *Let S be a scheme over $\text{Spec } \mathbb{Q}$ and G/S a reductive group scheme. Let $0 \rightarrow F_1 \rightarrow F_2 \rightarrow F_3 \rightarrow 0$ be an exact sequence of quasi coherent G -sheaves on S . Then the sequence*

$$0 \rightarrow F_1^G \rightarrow F_2^G \xrightarrow{v} F_3^G \rightarrow 0$$

is exact.

Proof. By (3.14.3.3) the above sequence is exact, except possibly at F_3^G . Thus we need to establish that v is surjective. By [Seshadri77, p.236] any quasi coherent G -sheaf is the union of its coherent G -subsheaves, thus it is sufficient to prove (7.3.3) for F_3 coherent. Then F_3^G is also coherent, hence, by the Nakayama lemma, v is surjective if it is surjective over closed points. This reduces (7.3.3) to the case when F_3 is a vector space over $k(s)$ for some closed point $s \in S$.

A similar problem, with S a scheme over $\text{Spec } \mathbb{Z}$, is considered in [Seshadri77, p.244], where it is proved that it reduces to the problem over $S = \text{Spec } \mathbb{Z}$. The same argument shows that if S is over $\text{Spec } \mathbb{Q}$, then (7.3.3) reduces to the special case where $S = \text{Spec } \mathbb{Q}$.

Over a field of characteristic zero any representation of a reductive group is completely reducible, thus (7.3.3) holds for $S = \text{Spec } \mathbb{Q}$. $\square$

7.4 Corollary. *Let G/S be a reductive group scheme and S a scheme over $\text{Spec } \mathbb{Q}$. Let $g : Y \rightarrow X$ be a G -morphism of algebraic spaces over S which is a closed immersion. Assume that the G -action on X is proper and the geometric quotient $f : X \rightarrow X/G$ exists.*

Then the geometric quotient $f : Y \rightarrow Y/G$ also exists and the induced morphism $f/G : Y/G \rightarrow X/G$ is a closed immersion.

Proof. The question is local in the étale topology of X/G , thus we may assume that X/G is affine. By (3.12.1) then X and Y are both affine. This is (7.3.2). $\square$

The main result of this section is the following

7.5 Theorem. *Let G be a reductive algebraic group scheme over S and X an algebraic space of finite type over S . Let $m : G \times X \rightarrow X$ be a proper G -action on X .*

Then the geometric quotient $f : X \rightarrow X/G$ exists and X/G is an algebraic space of finite type over S .

Proof. We already proved this result in case X is normal (4.2). Here we reduce the general case to the normal case.

By Noetherian induction we may assume that the geometric quotient exists for any G -invariant subscheme $Y \subset X$ (assuming $Y \neq X$).

First we consider the weakly normal case, and then the general case.

7.6 Proof of (7.5) in case X is weakly normal. We use the notation of (5.8). We need to check that the two conditions (5.8.1–2) are satisfied.

The geometric quotient $\bar{X}/G$ exists by (4.2) and C_X/G exists by the inductive assumption. Thus (5.8.1) is satisfied.

Next look at the diagram (5.8.2.1)

$$\begin{array}{ccc} Z(C_{\bar{X}}) & \xrightarrow{q_1} & Z(\bar{C}) \\ q_2 \downarrow & & \\ Z(C_X) & & \end{array}$$

$C_{\bar{X}} \rightarrow \bar{X}$ is a closed immersion, thus the induced morphism $Z(C_{\bar{X}}) \rightarrow Z(\bar{X})$ is a closed immersion at all points of characteristic zero by (7.4). $Z(\bar{C})$ is exactly the image of this morphism, thus $Z(C_{\bar{X}}) \xrightarrow{q_1} Z(\bar{C})$ is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

By (8.9) the above diagram can be completed to

$$\begin{array}{ccc} Z(C_{\bar{X}}) & \xrightarrow{q_1} & Z(\bar{C}) \\ q_2 \downarrow & & \downarrow s_2 \\ Z(C_X) & \xrightarrow{s_1} & W. \end{array}$$

Therefore, by (5.8), X/G exists. $\square$

7.7 Proof of (7.5) for arbitrary X . Let $n : X^{\text{wn}} \rightarrow X$ be the weak normalization. By (5.5), n is a finite and universal homeomorphism.

Let $f' : X^{\text{wn}} \rightarrow X^{\text{wn}}/G$ be the geometric quotient. Let U' be an affine scheme and $u : U' \rightarrow X^{\text{wn}}/G$ a surjective étale morphism. Set

$$V' = X^{\text{wn}} \times_{X^{\text{wn}}/G} U'.$$

f' is affine by (3.12.1), thus V' is also affine. The first projection $v' : V' \rightarrow X^{\text{wn}}$ is an étale and fixed-point reflecting G -morphism, since it is the pull back of the étale morphism u (2.13).

By (7.8) the fixed-point reflecting, surjective and étale G -morphism $V' \rightarrow X^{\text{wn}}$ descends to a fixed-point reflecting, surjective and étale G -morphism $V \rightarrow X$. $V' \rightarrow V$ is finite, hence V is also affine. The geometric quotient V/G exists by (7.3.1). Thus by (2.17) the geometric quotient X/G also exists. $\square$

7.8 Lemma. *Let G/S be a group scheme and $g : X' \rightarrow X$ a G -morphism which is a finite and universal homeomorphism. Then $V \mapsto V \times_X X'$ induces an equivalence between the categories*

$$(\text{étale } G\text{-morphisms: } * \rightarrow X) \xrightarrow{V \mapsto V \times_X X'} (\text{étale } G\text{-morphisms: } * \rightarrow X').$$

Moreover, $V \rightarrow X$ is fixed-point reflecting iff $V \times_X X' \rightarrow X'$ is.

Proof. Let $V' \rightarrow X'$ be an étale G -morphism. By (5.3) there is an étale morphism $V \rightarrow X$ such that $V' = V \times_X X'$. First we need to prove that G acts on V in the expected way. A G -action on V' is given by a morphism $m' : G \times V' \rightarrow V'$. We can encode this information as an isomorphism

$$M' : G \times V' \xrightarrow{(id_G, m')} G \times V'.$$

$G \times V' \rightarrow G \times X'$ is étale, thus by (5.3) M' descends to an isomorphism

$$M : G \times V \xrightarrow{(id_G, m)} G \times V.$$

The fact that m' is a G -action is expressed by certain commutative diagrams, which can be made to be diagrams of schemes étale over $G \times G \times X'$. By (5.3) these diagrams descend to X , showing that V is a G -scheme and $V' \rightarrow V$ is a G -morphism.

Finally, pick a point $v' \in V'$ and let $v \in V$, $x' \in X'$ and $x \in X$ denote its images. Since $V' \rightarrow V$ and $X' \rightarrow X$ are universal homeomorphisms, we see that

$$\begin{aligned} \text{red Stab}(v') &= \text{red Stab}(v) \quad \text{and} \quad \text{red Stab}(x') = \text{red Stab}(x), \quad \text{hence} \\ \text{red}(\text{Stab}(x')/\text{Stab}(v')) &= \text{red}(\text{Stab}(x)/\text{Stab}(v)). \end{aligned}$$

The fiber of $V' \rightarrow X'$ (resp. of $V \rightarrow X$) over x' (resp over x) is $\text{Stab}(x')/\text{Stab}(v')$ (resp. $\text{Stab}(x)/\text{Stab}(v)$). Both of these morphisms are étale, thus the fibers are smooth. They have the same reduced scheme structure, hence $\text{Stab}(x')/\text{Stab}(v') = \text{Stab}(x)/\text{Stab}(v)$. This implies that

$$\text{Stab}(v') = \text{Stab}(x') \quad \text{iff} \quad \text{Stab}(v) = \text{Stab}(x).$$

This completes the proof of (7.8). $\square$

8. PUSH-OUT DIAGRAMS

In this section we study push-out diagrams for algebraic spaces. These diagrams are used in going from normal to weakly-normal spaces.

8.1 Definition of Push-out Diagrams. The general problem is the following. Given a diagram

$$(8.1.1) \quad \begin{array}{ccc} X & \xrightarrow{g_1} & Y_1 \\ g_2 \downarrow & & \end{array}$$

we would like to complete it to a commutative diagram

$$(8.1.2) \quad \begin{array}{ccc} X & \xrightarrow{g_1} & Y_1 \\ g_2 \downarrow & & \downarrow s_1 \\ Y_2 & \xrightarrow{s_2} & Z. \end{array}$$

In the category of algebraic spaces over S this is always possible, take for instance $Z = S$. We are interested in finding an algebraic space Z which is “large”.

In all the cases that we consider, g_1 and g_2 are finite, and we would like s_1 and s_2 to be finite as well. In such a case we say that the push-out diagram has *finite morphisms*.

Assume that we have a diagram (8.1.2) with finite morphisms. Then there is a *universal push-out*, given by

$$(8.1.3) \quad Z^u := \operatorname{Spec}_Z \ker[\mathcal{O}_{Y_1} + \mathcal{O}_{Y_2} \xrightarrow{g_1^* - g_2^*} \mathcal{O}_X].$$

This implies that the construction of universal push-out diagrams is a local question over Z :

8.2 Lemma. *Assume that (8.1.2) is a universal push-out diagram with finite morphisms. Let $Z' \rightarrow Z$ be a flat morphism. Then the diagram obtained by base change*

$$\begin{array}{ccc} X \times_Z Z' & \xrightarrow{g'_1} & Y_1 \times_Z Z' \\ g'_2 \downarrow & & \downarrow s'_1 \\ Y_2 \times_Z Z' & \xrightarrow{s'_2} & Z' \end{array}$$

is again a universal push-out diagram with finite morphisms.

Thus the construction of universal push-out diagrams with finite morphisms is a local question in the étale topology of Z . $\square$

Unfortunately, in many cases finite push-out diagrams do not exist. The following examples give a sample of push-out diagrams with bad properties.

8.3 Examples. Let $S = \operatorname{Spec} k$ for a field k .

(8.3.1) Consider the diagram

$$\begin{array}{ccc} \mathbb{A}^1 & \xrightarrow{z^2} & \mathbb{A}^1 \\ z^2+z \downarrow & & \\ \mathbb{A}^1 & & \end{array}$$

The only possible push-out is $Z = \operatorname{Spec} k$, unless $\operatorname{char} k = 2$.

(8.3.2) (cf. [Holmann63, p. 342]) Let $X = \operatorname{Spec} k[x, y]$, $Y_1 = \operatorname{Spec} k[x, y^2, y^3]$ and $Y_2 = \operatorname{Spec} k[x + y, y^2, y^3]$. g_i are the natural morphisms; both are finite and universal homeomorphisms.

If $\operatorname{char} k = 0$ then the universal push-out is given by

$$Z = \operatorname{Spec}(k[x, y^2, y^3] \cap k[x + y, y^2, y^3]) = \operatorname{Spec} k[x^2, y^2, y^3, (x + y)^2, (x + y)^3]$$

which is not of finite type. The induced morphisms s_i are not finite.

The dependence on the characteristic becomes clear if we note that, as k -vector spaces

$$\begin{aligned} k[x, y^2, y^3] &= k[x^i y^2, x^i y^3 : i = 1, 2, \dots] + \sum_{i>0} kx^i, \quad \text{and} \\ k[x + y, y^2, y^3] &= k[x^i y^2, x^i y^3 : i = 1, 2, \dots] + \sum_{i>0} k(x^i + ix^{i-1}y). \end{aligned}$$

The second summands are disjoint in characteristic zero but have large intersection in positive characteristic.

(8.3.3) In (8.3.2) we may replace the polynomial rings with the corresponding power series rings. This shows that even in the category of algebraic or analytic spaces (in characteristic zero), there is no push-out with s_i finite.

In positive characteristic the situation is much better:

8.4 Lemma. *In the push-out diagram (8.1.1) assume that all algebraic spaces are of finite type over a scheme of positive characteristic. Assume furthermore that g_1 is a finite and universal homeomorphism. (No assumption on g_2 .) Then there is a push-out diagram*

$$\begin{array}{ccc} X & \xrightarrow{g_1} & Y_1 \\ g_2 \downarrow & & \downarrow s_1 \\ Y_2 & \xrightarrow{s_2} & Z, \end{array}$$

where s_2 is also a finite and universal homeomorphism.

Proof. The proof relies on properties of the Frobenius morphism (6.2). Choose q such that F^q factors as

$$F^q : X \xrightarrow{g_1} Y_1 \xrightarrow{\bar{g}_1} X^{(q)}.$$

Set $Z = Y_2^{(q)}$, $s_2 := F^q : Y_2 \rightarrow Y_2^{(q)}$ and $s_1 := g_2^{(q)} \circ \bar{g}_1$. $\square$

8.5 Example. In the situation of (8.4) the universal push-out may be rather complicated if g_2 is not finite. Consider for instance the diagram

$$\begin{array}{ccc} \operatorname{Spec} k[x] & \longrightarrow & \operatorname{Spec} k[x, y]/(y^2) \\ \downarrow & & \\ \operatorname{Spec} k & & \end{array}.$$

The construction of (8.4) gives $Z = \operatorname{Spec} k$. The universal push-out exists and is given by $Z^u = \operatorname{Spec}_k k[y, xy, x^2y, \dots]/(y^2)$. Z^u is a non-Noetherian scheme.

The most useful general result about push-out diagrams with finite morphisms is the following. As usual, we assume that the base scheme is excellent. The original more restrictive assumptions in [Artin70] can be relaxed in view of recent progress in the approximation property of Henselian rings [Popescu85; André94; Spisakowicz04].

8.6 Theorem. [Artin70, 6.1] *In the push-out diagram (8.1.1) assume that g_1 is a closed immersion and g_2 is finite. Then there is a push-out diagram*

$$\begin{array}{ccc} X & \xrightarrow{g_1} & Y_1 \\ g_2 \downarrow & & \downarrow s_1 \\ Y_2 & \xrightarrow{s_2} & Z, \end{array}$$

where s_2 is a closed immersion and s_1 is finite. $\square$

As (8.3.2) shows, the analog of (8.4) fails in characteristic zero. In order to get a general push-out result, we concentrate on finite and universal homeomorphisms which are isomorphisms in characteristic zero. First we give a ring theoretic characterization:

8.7 Lemma. *Let B be a Noetherian ring and $B_1 \subset B$ a subring such that B is finite over B_1 . The following are equivalent:*

(8.7.1) *$\text{Spec } B \rightarrow \text{Spec } B_1$ is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.*

(8.7.2) *There is an $m > 0$ such that $b^m \in B_1$ for every $b \in B$.*

Proof. Assume (8.7.2) and let $P < B_1$ be any prime ideal. Then B/PB is an Artin algebra over B_1/P such that if $b \in B/PB$ then $b^m \in B_1/P$. This implies that B/PB is local with maximal ideal $\sqrt{PB}$ and $B_1/P \subset B/\sqrt{PB}$ is a purely inseparable field extension. Thus by (5.2), $\text{Spec } B \rightarrow \text{Spec } B_1$ is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

Conversely, assume (8.7.1). B_1 is also Noetherian by (3.14.1) and B/B_1 is a finite B_1 -module. By assumption all associated primes of B/B_1 have positive residue characteristic, thus there is an integer N such that $NB \subset B_1$. We can replace the inclusion $B_1 \subset B$ by the inclusion $B_1/NB \subset B/NB$. Thus it is sufficient to prove (8.7) in case all the rings are $\mathbb{Z}/(N)$ -algebras.

B is a direct sum of $\mathbb{Z}/(p_i^{n_i})$ -algebras B^i where $N = \prod p_i^{n_i}$. Correspondingly, B_1 decomposes as a sum of subalgebras $B_1^i \subset B^i$.

Assume that for each i we find m_i such that $b^{m_i} \in B_1^i$ for every $b \in B^i$. Then $m = \prod m_i$ works for B . Thus we are further reduced to the case when all the rings are $\mathbb{Z}/(p^n)$ -algebras.

First consider $B_1/(B_1 \cap pB) \subset B/pB$. The induced map on the spectra is a finite and universal homeomorphism, hence by (6.6) there is an r such that $b^{p^r} \in B_1 + pB$ for every $b \in B$.

Let $b_1 + pb_2$ be a typical element of $B_1 + pB$. Then

$$(b_1 + pb_2)^{p^n} = b_1^{p^n} + \sum_{i>0} p^i \binom{p^n}{i} b_1^{p^n-i} b_2^i.$$

It is easy to see that p^n divides $p^i \binom{p^n}{i}$ for $i > 0$, thus $(b_1 + pb_2)^{p^n} = b_1^{p^n} \in B_1$. $\square$

8.8 Corollary. (8.8.1) *Assume that (8.1.2) is a universal push-out diagram with finite morphisms such that g_1 is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero. Then*

s_2 is also a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

(8.8.2) Assume that (8.1.1) is a push-out diagram with finite morphisms such that g_1 is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero. Then the construction of the universal push-out diagram is a local problem in the étale topology of Y_2 .

Proof. By (8.2) it is sufficient to consider (8.8.1) in case Z is an affine scheme. Then X and the Y_i are also affine. By (8.7) there is an $m > 0$ such that

$$b \in H^0(X, \mathcal{O}_X) \Rightarrow b^m \in \text{im} \left[H^0(Y_1, \mathcal{O}_{Y_1}) \xrightarrow{g_1^*} H^0(X, \mathcal{O}_X) \right].$$

Thus, by the construction (8.1.3) of the universal push-out,

$$b \in H^0(Y_2, \mathcal{O}_{Y_2}) \Rightarrow b^m \in \text{im} \left[H^0(Z, \mathcal{O}_Z) \xrightarrow{s_2^*} H^0(Y_2, \mathcal{O}_{Y_2}) \right].$$

By (8.7) this implies that s_2 is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

In order to see (8.8.2), let $V \rightarrow Y_2$ be an étale morphism. Since s_2 is a finite and universal homeomorphism, V descends to an étale cover $V_Z \rightarrow Z$ such that $V = V_Z \times_Z Y_2$ (5.4). Thus (8.2) and (8.8.1) imply (8.8.2). $\square$

This enables us to generalize (8.4) to algebraic spaces over an excellent base scheme S :

8.9 Lemma. *In the push-out diagram*

$$\begin{array}{ccc} X & \xrightarrow{g_1} & Y_1 \\ g_2 \downarrow & & \\ Y_2 & & \end{array}$$

assume that all algebraic spaces are of finite type over S . Assume that g_2 is finite and g_1 is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

Then there is a push-out diagram

$$\begin{array}{ccc} X & \xrightarrow{g_1} & Y_1 \\ g_2 \downarrow & & \downarrow s_1 \\ Y_2 & \xrightarrow{s_2} & Z, \end{array}$$

where s_1 is finite and s_2 is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

Proof. g_1 can be factored as

$$g_1 = h_1 \circ g'_1$$

where g'_1 is a closed immersion and $\mathcal{O}_{X_1} \rightarrow (h_1)_* \mathcal{O}_X$ is injective. We construct the push-out in two steps:

$$\begin{array}{ccccc} X & \xrightarrow{h_1} & X_1 & \xrightarrow{g'_1} & Y_1 \\ g_2 \downarrow & & \downarrow s'_1 & & \downarrow s_1 \\ Y_2 & \xrightarrow{s_2} & W & \xrightarrow{s'_2} & Z. \end{array}$$

By (8.8.2) the construction of W is local in the étale topology of Y_2 . Let $\text{Spec } C \subset S$ be an affine open set. Let U be an affine scheme of finite type over $\text{Spec } C$ and $U \rightarrow Y_2$ an étale morphism. Let $V = U \times_{Y_2} X$. Then $V \rightarrow U$ is finite, hence V is affine. Also, $V \rightarrow X$ is an étale morphism. Since g_1 is a universal homeomorphism, V descends to an affine scheme V_1 which is étale over X_1 (5.4). Corresponding to the morphisms of affine schemes $U \leftarrow V \rightarrow V_1$ we obtain morphisms of rings

$$A \xrightarrow{g_2^*} B \xleftarrow{h_1^*} B_1.$$

$B_1 \subset B$ is a subring such that $\text{Spec } B \rightarrow \text{Spec } B_1$ is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero. Choose m as in (8.7.2). Let $A_2 \subset A$ be the C -subalgebra generated by the m -th powers of all elements of A . A is a finitely generated C -algebra, thus A is finite over A_2 . By (8.7), $g_2(A_2) \subset B_1$. Set

$$A_1 := \{a \in A \mid g_2(a) \in B_1\}, \quad \text{and} \quad U_1 := \text{Spec } A_1.$$

Thus we obtain a commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{g_1} & V_1 \\ g_2 \downarrow & & \downarrow s_1 \\ U & \xrightarrow{s_2} & U_1, \end{array}$$

which gives the universal push-out. By (8.2.3) the schemes U_1 give an étale covering of W .

Once we have W , we can apply (8.6) to the diagram

$$\begin{array}{ccc} X_1 & \xrightarrow{g'_1} & Y_1 \\ \downarrow s'_1 & & \\ W & & \end{array}$$

to obtain Z . $\square$

In some applications (8.9) is used in the following slightly more general form:

8.10 Lemma. *In the push-out diagram*

$$\begin{array}{ccccc} X & \xrightarrow{g_1} & X_1 & \xrightarrow{h_1} & Y_1 \\ g_2 \downarrow & & & & \\ & & & & \end{array}$$

assume that g_2 is finite and h_1 is a closed immersion. Assume furthermore that g_1 is a finite and universal homeomorphism which is an isomorphism at all points where the characteristic of the residue field is zero.

Then there is a push-out diagram

$$\begin{array}{ccccc} X & \xrightarrow{g_1} & X_1 & \xrightarrow{h_1} & Y_1 \\ g_2 \downarrow & & & & \downarrow s_1 \\ Y_2 & \longrightarrow & & \longrightarrow & Z, \end{array}$$

where all morphisms are finite.

Proof. By (8.9) there is a push-out diagram

$$\begin{array}{ccc} X & \xrightarrow{g_1} & X_1 \\ g_2 \downarrow & & \downarrow h_2 \\ Y_2 & \longrightarrow & Y'_2, \end{array}$$

where all morphisms are finite. Then we can apply (8.6) to the diagram

$$\begin{array}{ccc} X_1 & \xrightarrow{h_1} & Y_1 \\ h_2 \downarrow & & \\ Y'_2 & & \end{array}$$

to find Z as required. $\square$

REFERENCES

- [André94] M. André, *Cinq exposés sur la désingularization (preprint)* (1994).
- [Andreotti-Bombieri69] A. Andreotti - E. Bombieri, *Sugli omeomorfismi delle varietà algebriche*, Ann. Sci. Norm. Sup. Pisa **23** (1969), 430-450.
- [Andreotti-Noguet67] A. Andreotti - F. Noguet, *La convexité holomorphe dans l'espace analytique des cycles d'une variété algébrique*, Ann. Sci. Norm. Sup. Pisa **21** (1967), 31-82.
- [Atiyah-MacDonald69] M. Atiyah - I. MacDonald, *Introduction to Commutative Algebra*, Addison-Wesley, 1969.
- [Artin69] M. Artin, *Algebraisation of formal moduli I.*, Global Analysis, D. C. Spenser and S. Iyanaga eds., Univ. Tokyo Press - Princeton Univ. Press, 1969, pp. 21-72.
- [Artin70] M. Artin, *Algebraisation of formal moduli II.*, Ann. of Math. **91** (1970), 88-135.
- [Artin74] M. Artin, *Versal deformations and algebraic stacks*, Inv. Math. **27** (1974), 165-189.
- [BLR90] S. Bosch - W. Lütkebohmert - M. Raynaud, *Néron Models*, Springer, 1990.
- [Demazure67] M. Demazure, *Schémas en groupes réductifs*, Bull. Soc. Math. France **93** (1967), 365-413.
- [Dethloff-Grauert94] G. Dethloff - H. Grauert, *Seminormal complex spaces*, Several Complex Variables, Enc. of Math. Sci. vol. VII., Springer, 1994, pp. 183-220.
- [Dixmier-Raynaud81] J. Dixmier - M. Raynaud, *Sur le quotient d'une variété algébrique*, Mathematical Analysis and Applications, L. Nachbin ed., Acad. Press, 1981, pp. 327-344.
- [EGA] A. Grothendieck - J. Dieudonné, *Eléments de Géométrie Algébrique*, vol. 4, 8, 11, 17, 20, 24, 28, 32, Publ. Math. IHES, 1960-67.
- [EGA71] A. Grothendieck - J. Dieudonné, *Eléments de Géométrie Algébrique I.*, Springer, 1971.
- [Fogarty83] J. Fogarty, *Geometric quotients are algebraic schemes*, Adv. in Math. **48** (1983), 166-171.

- [Greuel-Pfister93] G.-M. Greuel – G. Pfister, *Geometric quotients of unipotent group actions*, Proc. London Math. Soc. **67** (1993), 75-105.
- [Holmann60] H. Holmann, *Quotientenräume komplexer Mannigfaltigkeiten*, Math. Ann. **139** (1960), 383-402.
- [Holmann61] H. Holmann, *Quotienten komplexer Räume*, Math. Ann. **142** (1961), 407-440.
- [Holmann63] H. Holmann, *Komplexe Räume mit komplexen Transformationsgruppen*, Math. Ann. **150** (1963), 327-360.
- [Kaup68] B. Kaup, *Äquivalenzrelationen auf allgemeinen komplexen Räumen*, Schriftenreihe Math. Inst. Univ. Münster **39** (1968).
- [Keel94] S. Keel, *Lecture at the 1994 Salt Lake City Summer School on Moduli*.
- [Knutson71] D. Knutson, *Algebraic spaces*, Springer Lecture Notes vol. 203, 1971.
- [Kollár85] J. Kollár, *Toward moduli of singular varieties*, Comp. Math. **56** (1985), 369-398.
- [Kollár90] J. Kollár, *Projectivity of Complete Moduli*, J. Diff. Geom. **32** (1990), 235-268.
- [Kollár95a] J. Kollár, *Algebraic groups acting on schemes (preprint)* (1995).
- [Kollár95b] J. Kollár, *Rational Curves on Algebraic Varieties*, (to appear), 1995.
- [Luna73] D. Luna, *Slices Étales*, Bull. Soc. Math. France, Mémoires **33** (1973), 81-105.
- [Matsumura80] H. Matsumura, *Commutative Algebra*, (second edition), Benjamin - Cummings, 1980.
- [Matsusaka72] T. Matsusaka, *Polarised varieties with a given Hilbert polynomial*, Amer. J. Math. **94** (1972), 1027-1077.
- [Mumford65] D. Mumford, *Geometric invariant theory*, Springer, 1965.
- [Mumford-Fogarty82] D. Mumford - J. Fogarty, *Geometric invariant theory*, 2nd edition, Springer, 1982.
- [Nagata59] M. Nagata, *On the fourteenth problem of Hilbert*, Amer. J. Math. **81** (1959), 766-772.
- [Nagata69] M. Nagata, *Some questions on rational actions of groups*, Algebraic geometry, Bombay, Oxford Univ. Press., 1969, pp. 323-334.
- [Neeman89] A. Neeman, *Analytic questions in geometric invariant theory*, Invariant Theory, Contemp. Math. vol. 88, 1989, pp. 11-24.
- [Popescu85] D. Popescu, *General Néron desingularization*, Nagoya Math. J. **100** (1985), 97-126; ... and approximation, Nagoya Math. J. **104** (1986), 85-115.
- [Popp73] H. Popp, *On moduli of algebraic varieties I.*, Inv. Math. **22** (1973), 1-40; *II.*, Comp. Math. **28** (1974), 51-81; *III.*, Comp. Math. **31** (1975), 237-258.
- [Roberts86] M. Roberts, *A note on coherent G -sheaves*, Math. Ann. **275** (1986), 573-582.
- [Seshadri63] C. S. Seshadri, *Some results on the Quotient Space by an Algebraic Group of Automorphisms*, Math. Ann. **149** (1963), 286-301.
- [Seshadri72] C. S. Seshadri, *Quotient Spaces modulo Reductive Algebraic Groups*, Ann. Math. **95** (1972), 511-556.
- [Seshadri77] C. S. Seshadri, *Geometric reductivity over an arbitrary base*, Adv. in Math. **26** (1977), 225-274.
- [Seshadri82] C. S. Seshadri, *Fibrés Vectoriels sur les Courbes Algébriques*, Astérisque vol. 96, Soc. Math. France, 1982.
- [SGA3] M. Demazure - A. Grothendieck, *Schémas en Groupes*, Springer Lecture Notes vols. 151,152,153, 1970.
- [SGA5] A. Grothendieck, *Cohomologie l -adique et fonctions L* , Springer Lecture Notes vol. 589, 1977.
- [Spivakovsky95] M. Spivakovsky, *Smoothing of ring homomorphisms ...* (1995).
- [Traverso70] C. Traverso, *Seminormality and Picard Group*, Ann. Sci. E.N.S. Pisa **24** (1970), 585-595.
- [Viehweg95] E. Viehweg, *Quasi-Projective Moduli of Polarized Manifolds*, Springer, 1995.
- [Weil62] A. Weil, *Foundations of algebraic geometry*, AMS, 1962.

Department of Mathematics,
 University of Utah,
 Salt Lake City UT 84112, USA
 kollar@math.utah.edu